

MTBP (K-20): sc-47173

BACKGROUND

p53 is a critical coordinator of a wide range of stress responses. To facilitate a rapid response to stress, p53 is produced constitutively, but is negatively regulated by MDM2. MTBP (also designated MDM2BP or MDM2 transformed 3T3 cell double minute 2, p53 binding protein (mouse) binding protein) is a growth regulator that modulates the activity of MDM2 towards itself and p53, and thereby contributes to MDM2-dependent p53 homeostasis in cells. Specifically, MTBP promotes MDM2-mediated ubiquitination and degradation of p53 and also MDM2 stabilization. MTBP transcript is most abundant in thymus, testis and ovary.

REFERENCES

1. Boyd, M.T., et al. 2000. A novel cellular protein (MTBP) binds to and is suppressed by MDM2. *J. Biol. Chem.* 275: 31883-31890.
2. Boyd, M.T., et al. 2000. Assignment of 8q24 by *in situ* hybridization. *Cytogenet. Cell Genet.* 90: 64-65.

CHROMOSOMAL LOCATION

Genetic locus: MTBP (human) mapping to 8q24.12; Mtbp (mouse) mapping to 15 D1.

SOURCE

MTBP (K-20) is an affinity purified goat polyclonal antibody raised against a peptide mapping near the C-terminus of MTBP of mouse origin.

PRODUCT

Each vial contains 200 µg IgG in 1.0 ml of PBS with < 0.1% sodium azide and 0.1% gelatin.

Blocking peptide available for competition studies, sc-47173 P, (100 µg peptide in 0.5 ml PBS containing < 0.1% sodium azide and 0.2% BSA).

APPLICATIONS

MTBP (K-20) is recommended for detection of MTBP of mouse and human origin by Western Blotting (starting dilution 1:200, dilution range 1:100-1:1000), immunoprecipitation [1-2 µg per 100-500 µg of total protein (1 ml of cell lysate)], immunofluorescence (starting dilution 1:50, dilution range 1:50-1:500) and solid phase ELISA (starting dilution 1:30, dilution range 1:30-1:3000).

MTBP (K-20) is also recommended for detection of MTBP in additional species, including equine, canine, bovine and porcine.

Suitable for use as control antibody for MTBP siRNA (m): sc-61081, MTBP siRNA (h): sc-61080, MTBP shRNA Plasmid (m): sc-61081-SH, MTBP shRNA Plasmid (h): sc-61080-SH, MTBP shRNA (m) Lentiviral Particles: sc-61081-V and MTBP shRNA (h) Lentiviral Particles: sc-61080-V.


Molecular Weight of MTBP: 104 kDa.

Positive Controls: MTBP (m): 293T Lysate: sc-121821, K-562 whole cell lysate: sc-2203 or 3T3-L1 cell lysate: sc-2243.


RECOMMENDED SECONDARY REAGENTS

To ensure optimal results, the following support (secondary) reagents are recommended: 1) Western Blotting: use donkey anti-goat IgG-HRP: sc-2020 (dilution range: 1:2000-1:100,000) or Cruz Marker™ compatible donkey anti-goat IgG-HRP: sc-2033 (dilution range: 1:2000-1:5000), Cruz Marker™ Molecular Weight Standards: sc-2035, TBS Blotto A Blocking Reagent: sc-2333 and Western Blotting Luminol Reagent: sc-2048. 2) Immunoprecipitation: use Protein A/G PLUS-Agarose: sc-2003 (0.5 ml agarose/2.0 ml). 3) Immunofluorescence: use donkey anti-goat IgG-FITC: sc-2024 (dilution range: 1:100-1:400) or donkey anti-goat IgG-TR: sc-2783 (dilution range: 1:100-1:400) with UltraCruz™ Mounting Medium: sc-24941.

DATA



MTBP (K-20): sc-47173. Western blot analysis of MTBP expression in non-transfected: sc-117752 (A) and mouse MTBP transfected: sc-121821 (B) 293T whole cell lysates.



MTBP (K-20): sc-47173. Western blot analysis of MTBP expression in 3T3-L1 whole cell lysate.

SELECT PRODUCT CITATIONS

1. Iwakuma, T., et al. 2008. Mtbp haploinsufficiency in mice increases tumor metastasis. *Oncogene* 27: 1813-1820.
2. Agarwal, N., et al. 2011. MTBP plays a crucial role in mitotic progression and chromosome segregation. *Cell Death Differ.* 18: 1208-1219.

STORAGE

Store at 4° C, **DO NOT FREEZE**. Stable for one year from the date of shipment. Non-hazardous. No MSDS required.

RESEARCH USE

For research use only, not for use in diagnostic procedures.

PROTOCOLS

See our web site at www.scbt.com or our catalog for detailed protocols and support products.

MONOS
Satisfaction
Guaranteed

Try **MTBP (B-5): sc-137201**, our highly recommended monoclonal alternative to MTBP (K-20).