

NAT-1/2 (K-14): sc-47230

BACKGROUND

Arylamine N-acetyltransferases (NAT-1 and NAT-2) catalyze N- or O-acetylation of heterocyclic and arylamine substrates in the detoxification of a wide array of drugs. Certain alleles causing high levels of N-acetyltransferase activity have been associated with colon and urinary bladder cancers, as NAT's also bioactivate several known carcinogens. Both NAT-1 and NAT-2 are cytoplasmic proteins and play an active role in the detoxification of many arylamine and hydrazine drugs. N-acetylation polymorphism is determined by the level of NAT activity in liver tissues, and has been linked to the action and toxicity of drugs that contain amines. Human NAT-1 is the functional homolog of rodent NAT-2, while human NAT-2 is the functional homolog of rodent NAT-1.

CHROMOSOMAL LOCATION

Genetic locus: NAT1/NAT2 (human) mapping to 8p22; Nat1/Nat2 (mouse) mapping to 8 B3.3.

SOURCE

NAT-1/2 (K-14) is an affinity purified goat polyclonal antibody raised against a peptide mapping within an internal region of NAT-2 of mouse origin.

PRODUCT

Each vial contains 200 µg IgG in 1.0 ml of PBS with < 0.1% sodium azide and 0.1% gelatin.

Blocking peptide available for competition studies, sc-47230 P, (100 µg peptide in 0.5 ml PBS containing < 0.1% sodium azide and 0.2% BSA).

STORAGE

Store at 4° C, ****DO NOT FREEZE****. Stable for one year from the date of shipment. Non-hazardous. No MSDS required.

APPLICATIONS

NAT-1/2 (K-14) is recommended for detection of NAT-1 of human origin and NAT-2 of mouse and rat origin by Western Blotting (starting dilution 1:200, dilution range 1:100-1:1000), immunoprecipitation [1-2 µg per 100-500 µg of total protein (1 ml of cell lysate)], immunofluorescence (starting dilution 1:50, dilution range 1:50-1:500) and solid phase ELISA (starting dilution 1:30, dilution range 1:30-1:3000).

NAT-1/2 (K-14) is also recommended for detection of NAT-1 and NAT-2 in additional species, including bovine, porcine and avian.

Suitable for use as control antibody for NAT-1 siRNA (h): sc-61154, NAT-1 shRNA Plasmid (h): sc-61154-SH, NAT-1 shRNA (h) Lentiviral Particles: sc-61154-V, NAT-2 siRNA (m): sc-61157, NAT-2 shRNA Plasmid (m): sc-61157-SH, and NAT-2 shRNA (m) Lentiviral Particles: sc-61157-V.


Molecular Weight of NAT-1/2: 34 kDa.

Positive Controls: NAT-2 (m): 293T Lysate: sc-121945 or HeLa whole cell lysate: sc-2200.

RECOMMENDED SECONDARY REAGENTS

To ensure optimal results, the following support (secondary) reagents are recommended: 1) Western Blotting: use donkey anti-goat IgG-HRP: sc-2020 (dilution range: 1:2000-1:100,000) or Cruz Marker™ compatible donkey anti-goat IgG-HRP: sc-2033 (dilution range: 1:2000-1:5000), Cruz Marker™ Molecular Weight Standards: sc-2035, TBS Blotto A Blocking Reagent: sc-2333 and Western Blotting Luminol Reagent: sc-2048. 2) Immunoprecipitation: use Protein A/G PLUS-Agarose: sc-2003 (0.5 ml agarose/2.0 ml). 3) Immunofluorescence: use donkey anti-goat IgG-FITC: sc-2024 (dilution range: 1:100-1:400) or donkey anti-goat IgG-TR: sc-2783 (dilution range: 1:100-1:400) with UltraCruz™ Mounting Medium: sc-24941.

DATA



NAT-1/2 (K-14): sc-47230. Western blot analysis of NAT-2 expression in non-transfected: sc-117752 (A) and mouse NAT-2 transfected: sc-121945 (B) 293T whole cell lysates.

RESEARCH USE

For research use only, not for use in diagnostic procedures.

PROTOCOLS

See our web site at www.scbt.com or our catalog for detailed protocols and support products.

MONOS
Satisfaction
Guaranteed

Try **NAT-1/2 (G-5): sc-137204** or **NAT-1/2 (H-7): sc-271797**, our highly recommended monoclonal alternatives to NAT-1/2 (K-14).