

Pancreatic Lipase (A-12): sc-47291

BACKGROUND

The lipase gene family belongs to one of the most robust genetic superfamilies found in living organisms, which includes esterases and thioesterases. Members of the AB hydrolase subfamily include Hepatic Lipase (HL), Endothelial lipase (EL), Lipoprotein Lipase (LPL), Pancreatic Lipase (PL), Gastric Lipase (GL) and LCAT. These family members play a crucial role in the metabolism of lipids. Pancreatic Lipase, also designated pancreatic triacylglycerol acyl hydrolase, is important for dietary fat absorption as it hydrolyses triglycerides into diglycerides, monoglycerides and free fatty acids.

REFERENCES

1. Lowe, M.E., Rosenblum, J.L. and Strauss, A.W. 1989. Cloning and characterization of human Pancreatic Lipase cDNA. *J. Biol. Chem.* 264: 20042-20048.
2. Winkler, F.K., D'Arcy, A. and Hunziker, W. 1990. Structure of human Pancreatic Lipase. *Nature* 343: 771-774.
3. Yajima, H., Noguchi, T., Ikeshima, E., Shiraki, M., Kanaya, T., Tsuboyama-Kasaoka, N., Ezaki, O., Oikawa, S. and Kondo, K. 2005. Prevention of diet-induced obesity by dietary isomerized hop extract containing isohumulones, in rodents. *Int. J. Obes. Relat. Metab. Disord.* 29:991-997.
4. Han, L.K., Zheng, Y.N., Yoshikawa, M., Okuda, H. and Kimura, Y. 2005. Anti-obesity effects of chikusetsusaponins isolated from *Panax japonicus* rhizomes. *BMC Complement Altern. Med.* 5:9.
5. Bijvelds, M.J., Bronsveld, I., Havinga, R., Sinaasappel, M., de Jonge, H.R. and Verkade, H.J. 2005. Fat absorption in cystic fibrosis mice is impeded by defective lipolysis and post-lipolytic events. *Am. J. Physiol. Gastrointest. Liver Physiol.* 288: G646-653.
6. Sharma, N., Sharma, V.K. and Seo, S.Y. 2005. Screening of some medicinal plants for anti-lipase activity. *J. Ethnopharmacol.* 97: 453-456.
7. Pappan, K.L., Pan, Z., Kwon, G., Marshall, C.A., Coleman, T., Goldberg, I.J., McDaniel, M.L. and Semenkovich, C.F. 2005. Pancreatic β cell lipoprotein lipase independently regulates islet glucose metabolism and normal insulin secretion. *J. Biol. Chem.* 280: 9023-9029.

CHROMOSOMAL LOCATION

Genetic locus: PNLIP (human) mapping to 10q26.1; Pnlip (mouse) mapping to 19 D2.

SOURCE

Pancreatic Lipase (A-12) is an affinity purified goat polyclonal antibody raised against a peptide mapping within an internal region of Pancreatic Lipase of human origin.

PRODUCT

Each vial contains 200 μ g IgG in 1.0 ml of PBS with < 0.1% sodium azide and 0.1% gelatin.

Blocking peptide available for competition studies, sc-47291 P, (100 μ g peptide in 0.5 ml PBS containing < 0.1% sodium azide and 0.2% BSA).

APPLICATIONS

Pancreatic Lipase (A-12) is recommended for detection of Pancreatic Lipase of human origin by Western Blotting (starting dilution 1:200, dilution range 1:100-1:1000), immunofluorescence (starting dilution 1:50, dilution range 1:50-1:500) and solid phase ELISA (starting dilution 1:30, dilution range 1:30-1:3000).

Suitable for use as control antibody for Pancreatic Lipase siRNA (h): sc-61285.

Molecular Weight of Pancreatic Lipase: 51 kDa.

RECOMMENDED SECONDARY REAGENTS

To ensure optimal results, the following support (secondary) reagents are recommended: 1) Western Blotting: use donkey anti-goat IgG-HRP: sc-2020 (dilution range: 1:2000-1:100,000) or Cruz Marker™ compatible donkey anti-goat IgG-HRP: sc-2033 (dilution range: 1:2000-1:5000), Cruz Marker™ Molecular Weight Standards: sc-2035, TBS Blotto A Blocking Reagent: sc-2333 and Western Blotting Luminol Reagent: sc-2048. 2) Immunofluorescence: use donkey anti-goat IgG-FITC: sc-2024 (dilution range: 1:100-1:400) or donkey anti-goat IgG-TR: sc-2783 (dilution range: 1:100-1:400) with UltraCruz™ Mounting Medium: sc-24941.

STORAGE

Store at 4° C, **DO NOT FREEZE**. Stable for one year from the date of shipment. Non-hazardous. No MSDS required.

RESEARCH USE

For research use only, not for use in diagnostic procedures.

PROTOCOLS

See our web site at www.scbt.com or our catalog for detailed protocols and support products.



Try **Pancreatic Lipase (A-3): sc-374612**, our highly recommended monoclonal alternative to Pancreatic Lipase (A-12).