

β-taxilin (C-18): sc-47456

BACKGROUND

α-taxilin is a novel binding partner of the syntaxin family, which is implicated in intracellular vesicle trafficking. Through its C-terminal coiled-coil region, α-taxilin interacts with the nascent polypeptide-associated complex (NAC), which acts as a transcriptional coactivator. Although α-taxilin binds to both the α and β NAC subunits, the main interaction is through αNAC. Co-expression of α-taxilin with overexpressed αNAC eliminated the nuclear distribution of αNAC, originally distributed throughout the cytosol and nucleus. In addition, other taxilin family members, β- and γ-taxilins, bind to αNAC and affect its nuclear distribution, suggesting that the taxilin family is involved not only in the translational process through its interaction with NAC but also in the transcriptional process through its interaction with αNAC alone.

REFERENCES

1. Nogami, S., et al. 2003. Interaction of taxilin with syntaxin which does not form the SNARE complex. *Biochem. Biophys. Res. Commun.* 311: 797-802.
2. Nogami, S., et al. 2003. Taxilin; a novel syntaxin-binding protein that is involved in Ca²⁺-dependent exocytosis in neuroendocrine cells. *Genes Cells* 8:17-28.
3. Nogami, S., et al. 2004. Identification and characterization of taxilin isoforms. *Biochem. Biophys. Res. Commun.* 319: 936-943.
4. Yoshida, K., et al. 2005. Interaction of the taxilin family with the nascent polypeptide-associated complex that is involved in the transcriptional and translational processes. *Genes Cells* 10: 465-476.
5. Malyala, A., et al. 2005. Estrogen modulation of hypothalamic neurons: Activation of multiple signaling pathways and gene expression changes. *Steroids* 70: 397-406.

CHROMOSOMAL LOCATION

Genetic locus: MDP77 (human) mapping to 6q24.1.

SOURCE

β-taxilin (C-18) is an affinity purified goat polyclonal antibody raised against a peptide mapping at the C-terminus of β-taxilin of human origin.

PRODUCT

Each vial contains 200 μg IgG in 1.0 ml of PBS with < 0.1% sodium azide and 0.1% gelatin.

Blocking peptide available for competition studies, sc-47456 P, (100 μg peptide in 0.5 ml PBS containing < 0.1% sodium azide and 0.2% BSA).

STORAGE

Store at 4° C, ****DO NOT FREEZE****. Stable for one year from the date of shipment. Non-hazardous. No MSDS required.

PROTOCOLS

See our web site at www.scbt.com or our catalog for detailed protocols and support products.

APPLICATIONS

β-taxilin (C-18) is recommended for detection of β-taxilin of human origin by Western Blotting (starting dilution 1:200, dilution range 1:100-1:1000), immunoprecipitation [1-2 μg per 100-500 μg of total protein (1 ml of cell lysate)], immunofluorescence (starting dilution 1:50, dilution range 1:50-1:500) and solid phase ELISA (starting dilution 1:30, dilution range 1:30-1:3000).

Suitable for use as control antibody for β-taxilin siRNA (h): sc-61651, β-taxilin shRNA Plasmid (h): sc-61651-SH and β-taxilin shRNA (h) Lentiviral Particles: sc-61651-V.


Molecular Weight of β-taxilin: 120 kDa.

Positive Controls: β-taxilin (h3): 293T Lysate: sc-129920.

RECOMMENDED SECONDARY REAGENTS

To ensure optimal results, the following support (secondary) reagents are recommended: 1) Western Blotting: use donkey anti-goat IgG-HRP: sc-2020 (dilution range: 1:2000-1:100,000) or Cruz Marker™ compatible donkey anti-goat IgG-HRP: sc-2033 (dilution range: 1:2000-1:5000), Cruz Marker™ Molecular Weight Standards: sc-2035, TBS Blotto A Blocking Reagent: sc-2333 and Western Blotting Luminol Reagent: sc-2048. 2) Immunoprecipitation: use Protein A/G PLUS-Agarose: sc-2003 (0.5 ml agarose/2.0 ml). 3) Immunofluorescence: use donkey anti-goat IgG-FITC: sc-2024 (dilution range: 1:100-1:400) or donkey anti-goat IgG-TR: sc-2783 (dilution range: 1:100-1:400) with UltraCruz™ Mounting Medium: sc-24941.

DATA



β-taxilin (C-18): sc-47456. Western blot analysis of β-taxilin expression in non-transfected: sc-117752 (A) and human β-taxilin transfected: sc-129920 (B) 293T whole cell lysates.

RESEARCH USE

For research use only, not for use in diagnostic procedures.