

ARP-1 (K-14): sc-47578

BACKGROUND

COUP (chicken ovalbumin upstream promoter) transcription factors have been cloned in several species and identified as orphan members of the steroid/thyroid hormone receptor superfamily. COUP-TFI (also designated COUP or EAR-3) and ARP-1 (also designated COUP-TFII) exhibit highly regulated and overlapping expression in most tissues. COUP-TFs are highly expressed in the developing and central nervous system, suggesting that these factors may be important in neural development and differentiation. COUP-TFs can compete for binding to response elements which are common to other members of this family, including RAR, RXR, PPAR, HNF-4, VDR and TR. They have been shown to act as negative regulators as well as initiators of transcription.

CHROMOSOMAL LOCATION

Genetic locus: NR2F2 (human) mapping to 15q26.2; Nr2f2 (mouse) mapping to 7 D1.

SOURCE

ARP-1 (K-14) is an affinity purified goat polyclonal antibody raised against a peptide mapping near the N-terminus of ARP-1 of human origin.

PRODUCT

Each vial contains 200 µg IgG in 1.0 ml of PBS with < 0.1% sodium azide and 0.1% gelatin. Also available as TransCruz reagent for Gel Supershift and ChIP applications, sc-47578 X, 200 µg/0.1 ml.

Blocking peptide available for competition studies, sc-47578 P, (100 µg peptide in 0.5 ml PBS containing < 0.1% sodium azide and 0.2% BSA).

RESEARCH USE

For research use only, not for use in diagnostic procedures.

APPLICATIONS

ARP-1 (K-14) is recommended for detection of ARP-1 of mouse, rat and human origin by Western Blotting (starting dilution 1:200, dilution range 1:100-1:1000), immunoprecipitation [1-2 µg per 100-500 µg of total protein (1 ml of cell lysate)], immunofluorescence (starting dilution 1:50, dilution range 1:50-1:500) and solid phase ELISA (starting dilution 1:30, dilution range 1:30-1:3000); may cross-react with COUP-TFI.

ARP-1 (K-14) is also recommended for detection of ARP-1 in additional species, including equine, canine, bovine, porcine and avian.

Suitable for use as control antibody for ARP-1 siRNA (h): sc-38818, ARP-1 siRNA (m): sc-38819, ARP-1 shRNA Plasmid (h): sc-38818-SH, ARP-1 shRNA Plasmid (m): sc-38819-SH, ARP-1 shRNA (h) Lentiviral Particles: sc-38818-V and ARP-1 shRNA (m) Lentiviral Particles: sc-38819-V.

ARP-1 (K-14) X TransCruz antibody is recommended for Gel Supershift and ChIP applications.


Molecular Weight of ARP-1: 45 kDa.

Positive Controls: ARP-1 (h): 293T Lysate: sc-111675 or Ramos nuclear extract: sc-2153.

RECOMMENDED SECONDARY REAGENTS

To ensure optimal results, the following support (secondary) reagents are recommended: 1) Western Blotting: use donkey anti-goat IgG-HRP: sc-2020 (dilution range: 1:2000-1:100,000) or Cruz Marker™ compatible donkey anti-goat IgG-HRP: sc-2033 (dilution range: 1:2000-1:5000), Cruz Marker™ Molecular Weight Standards: sc-2035, TBS Blotto A Blocking Reagent: sc-2333 and Western Blotting Luminol Reagent: sc-2048. 2) Immunofluorescence: use donkey anti-goat IgG-FITC: sc-2024 (dilution range: 1:100-1:400) or donkey anti-goat IgG-TR: sc-2783 (dilution range: 1:100-1:400) with UltraCruz™ Mounting Medium: sc-24941.

DATA



ARP-1 (K-14): sc-47578. Western blot analysis of ARP-1 expression in non-transfected: sc-117752 (A) and human ARP-1 transfected: sc-111675 (B) 293T whole cell lysates.

STORAGE

Store at 4° C, ****DO NOT FREEZE****. Stable for one year from the date of shipment. Non-hazardous. No MSDS required.

PROTOCOLS

See our web site at www.scbt.com or our catalog for detailed protocols and support products.


 MONOS
Satisfaction
Guaranteed

Try **ARP-1 (E-11): sc-271940** or **ARP-1 (B-2): sc-271265**, our highly recommended monoclonal alternatives to ARP-1 (K-14).