



NEIL2 (G-13): sc-47613

BACKGROUND

NEIL1, NEIL2 and NEIL3, also known as endonuclease VIII-like 1, 2 and 3 or DNA-(apurinic or apyrimidinic site) lyase NEIL 1, 2 and 3, are nuclear proteins involved in the repair of DNA damaged by oxidation. The NEIL proteins belong to the FPG family of proteins. They act as DNA glycosylases that can recognize and remove damaged bases, leaving an abasic site. NEIL3, however, lacks the proline residue at the N-terminus which acts as the active site residue found in NEIL1 and NEIL2. NEIL1 is a ubiquitously expressed protein that is upregulated during S phase. NEIL2 is expressed primarily in testis, heart, skeletal muscle, placenta, brain, kidney and liver while NEIL3 is detected primarily in thymus and testis.

REFERENCES

- Hazra, T.K., Kow, Y.W., Hatahet, Z., Imhoff, B., Boldogh, I., Mokkalapati, S.K., Mitra, S. and Izumi, T. 2002. Identification and characterization of a novel human DNA glycosylase for repair of cytosine-derived lesions. *J. Biol. Chem.* 277: 30417-30420.
- Dou, H., Mitra, S. and Hazra, T.K. 2003. Repair of oxidized bases in DNA bubble structures by human DNA glycosylases NEIL1 and NEIL2. *J. Biol. Chem.* 278: 49679-49684.
- Shimura, K., Tao, H., Goto, M., Igarashi, H., Taniguchi, T., Maekawa, M., Takezaki, T. and Sugimura, H. 2004. Inactivating mutations of the human base excision repair gene NEIL1 in gastric cancer. *Carcinogenesis* 25: 2311-2317.
- Das, A., Rajagopalan, L., Mathura, V.S., Rigby, S.J., Mitra, S. and Hazra, T.K. 2004. Identification of a zinc finger domain in the human NEIL2 protein. *J. Biol. Chem.* 279: 47132-47138.
- Mokkalapati, S.K., Wiederhold, L., Hazra, T.K. and Mitra, S. 2004. Stimulation of DNA glycosylase activity of OGG1 by NEIL1: functional collaboration between two human DNA glycosylases. *Biochemistry* 43: 11596-11604.
- Hailer, M.K., Slade, P.G., Martin, B.D., Rosenquist, T.A. and Sugden, K.D. 2005. Recognition of the oxidized lesions spiroiminodihydroantoin and guanidinohydroantoin in DNA by the mammalian base excision repair DNA. *DNA Repair* 4: 41-50.

CHROMOSOMAL LOCATION

Genetic locus: (human) mapping to ; (mouse) mapping to .

SOURCE

NEIL2 (G-13) is an affinity purified goat polyclonal antibody raised against a peptide mapping near the C-terminus of NEIL2 of human origin.

PRODUCT

Each vial contains 200 µg IgG in 1.0 ml of PBS with < 0.1% sodium azide and 0.1% gelatin.

Blocking peptide available for competition studies, sc-47613 P, (100 µg peptide in 0.5 ml PBS containing < 0.1% sodium azide and 0.2% BSA).

Available as TransCruz reagent for Gel Supershift and ChIP applications, sc-47613 X, 200 µg/0.1 ml.

APPLICATIONS

NEIL2 (G-13) is recommended for detection of NEIL2 of human origin by Western Blotting (starting dilution 1:200, dilution range 1:100-1:1000) and immunofluorescence (starting dilution 1:50, dilution range 1:50-1:500).

Suitable for use as control antibody for NEIL2 siRNA (h): sc-61168.

NEIL2 (G-13) X TransCruz antibody is recommended for Gel Supershift and ChIP applications.

RECOMMENDED SECONDARY REAGENTS

To ensure optimal results, the following support (secondary) reagents are recommended: 1) Western Blotting: use donkey anti-goat IgG-HRP: sc-2020 (dilution range: 1:2000-1:100,000) or Cruz Marker™ compatible donkey anti-goat IgG-HRP: sc-2033 (dilution range: 1:2000-1:5000), Cruz Marker™ Molecular Weight Standards: sc-2035, TBS Blotto A Blocking Reagent: sc-2333 and Western Blotting Luminol Reagent: sc-2048. 2) Immunofluorescence: use donkey anti-goat IgG-FITC: sc-2024 (dilution range: 1:100-1:400) or donkey anti-goat IgG-TR: sc-2783 (dilution range: 1:100-1:400) with UltraCruz™ Mounting Medium: sc-24941.

STORAGE

Store at 4° C, ****DO NOT FREEZE****. Stable for one year from the date of shipment. Non-hazardous. No MSDS required.

RESEARCH USE

For research use only, not for use in diagnostic procedures.

PROTOCOLS

See our web site at www.scbt.com or our catalog for detailed protocols and support products.