

CYP11B1 (V-12): sc-47652

BACKGROUND

The steroid 11 β -hydroxylase gene, also designated CYP11B1, is a marker for the functional differentiation of cells in the zonae fasciculata reticularis. The deduced protein CYP11B1 consists of 466 amino acids containing a secretory signal, epidermal growth factor-like repeats, and a proteolytically inactive cathepsin B-related sequence. A related protein, human aldosterone synthase (CYP11B2), is involved in substrate recognition and conversion, with a functionally significant residue 112 in the N-terminal region of human CYP11B2. The inherited disorder glucocorticoid-remediable aldosteronism is caused by a chimeric gene duplication between the CYP11B1 and CYP11B2 genes. This disorder is characterized by hyperaldosteronism and high levels of 18-hydroxy-cortisol and 18-oxocortisol, which are under ACTH control.

REFERENCES

- Fardella, C.E., et al. 2001. Genetic study of patients with dexamethasone-suppressible aldosteronism without the chimeric CYP11B1/CYP11B2 gene. *J. Clin. Endocrinol. Metab.* 86: 4805-4807.
- Bechtel, S., et al. 2002. The effect of amino-acid substitutions I112P, D147E and K152N in CYP11B2 on the catalytic activities of the enzyme. *Eur. J. Biochem.* 269: 1118-1127.
- Mukai, K., et al. 2003. An inverse correlation between expression of a pre-cathepsin B-related protein with cysteine-rich sequences and steroid 11 β -hydroxylase in adrenocortical cells. *J. Biol. Chem.* 278: 17084-17092.
- Ganapathipillai, S., et al. 2005. CYP11B2-CYP11B1 haplotypes associated with decreased 11 β -hydroxylase activity. *J. Clin. Endocrinol. Metab.* 90: 1220-1225.
- Krone, N., et al. 2005. Congenital adrenal hyperplasia due to 11-hydroxylase deficiency: functional characterization of two novel point mutations and a three-base pair deletion in the CYP11B1 gene. *J. Clin. Endocrinol. Metab.* 90: 3724-3730.
- Barr, M., et al. 2006. Functional effects of genetic variants in the 11 β -hydroxylase (CYP11B1) gene. *Clin. Endocrinol.* 65: 816-825.
- Hakki, T., et al. 2008. Coexpression of redox partners increases the hydrocortisone (cortisol) production efficiency in CYP11B1 expressing fission yeast *Schizosaccharomyces pombe*. *J. Biotechnol.* 133: 351-359.

CHROMOSOMAL LOCATION

Genetic locus: Cyp11b1 (rat) mapping to 7q34.

SOURCE

CYP11B1 (V-12) is an affinity purified goat polyclonal antibody raised against a peptide mapping within an internal region of CYP11B1 of rat origin.

PRODUCT

Each vial contains 200 μ g IgG in 1.0 ml of PBS with < 0.1% sodium azide and 0.1% gelatin.

Blocking peptide available for competition studies, sc-47652 P, (100 μ g peptide in 0.5 ml PBS containing < 0.1% sodium azide and 0.2% BSA).

APPLICATIONS

CYP11B1 (V-12) is recommended for detection of CYP11B1 of rat origin by Western Blotting (starting dilution 1:200, dilution range 1:100-1:1000), immunoprecipitation [1-2 μ g per 100-500 μ g of total protein (1 ml of cell lysate)], immunofluorescence (starting dilution 1:50, dilution range 1:50-1:500) and solid phase ELISA (starting dilution 1:30, dilution range 1:30-1:3000).


Molecular Weight of CYP11B1: 48 kDa.

Positive Controls: rat adrenal gland extract: sc-364802.

RECOMMENDED SECONDARY REAGENTS

To ensure optimal results, the following support (secondary) reagents are recommended: 1) Western Blotting: use donkey anti-goat IgG-HRP: sc-2020 (dilution range: 1:2000-1:100,000) or Cruz Marker™ compatible donkey anti-goat IgG-HRP: sc-2033 (dilution range: 1:2000-1:5000), Cruz Marker™ Molecular Weight Standards: sc-2035, TBS Blotto A Blocking Reagent: sc-2333 and Western Blotting Luminol Reagent: sc-2048. 2) Immunoprecipitation: use Protein A/G PLUS-Agarose: sc-2003 (0.5 ml agarose/2.0 ml). 3) Immunofluorescence: use donkey anti-goat IgG-FITC: sc-2024 (dilution range: 1:100-1:400) or donkey anti-goat IgG-TR: sc-2783 (dilution range: 1:100-1:400) with UltraCruz™ Mounting Medium: sc-24941.

DATA



CYP11B1 (V-12): sc-47652. Western blot analysis of CYP11B1 expression in rat adrenal gland tissue extract.

STORAGE

Store at 4° C, **DO NOT FREEZE**. Stable for one year from the date of shipment. Non-hazardous. No MSDS required.

RESEARCH USE

For research use only, not for use in diagnostic procedures.



Try **CYP11B1 (A-11): sc-377401** or **CYP11B1 (B-11): sc-377248**, our highly recommended monoclonal alternatives to CYP11B1 (V-12).