

Met/Leu-enkephalin (NOC1/35): sc-47705

BACKGROUND

The corticotropin-lipotropin protein, also designated pro-opiomelanocortin, belongs to the POMC family of proteins. Specific cleavages by enzymes at certain basic paired residues yield various processed active peptides. The corticotropin-lipotropin protein contains synenkephalin, Leu-enkephalin, NPP, α -, β - and γ -melanotropin (MSH), adrenocorticotrophic hormone (ACTH), lipotropin β and γ , β -endorphin and Met-enkephalin processed active peptides. β -endorphin and Met-enkephalin are endogenous opiates while ACTH is crucial for adrenal gland stimulation to release cortisol. Met/Leu-enkephalins also function in pain perception and response to stress. MSH increased melanin production in melanocytes which leads to an increase in skin pigmentation. Leumorphin may be important in apoptosis prevention by being involved in the MAP-K and PI3-K pathways.

CHROMOSOMAL LOCATION

Genetic locus: PENK (human) mapping to 8q12.1, POMC (human) mapping to 2p23.3; Penk (mouse) mapping to 4 A1, Pomc (mouse) mapping to 12 A1.1.

SOURCE

Met/Leu-enkephalin (NOC1/35) is a mouse monoclonal antibody raised against a synthetic peptide corresponding to Leu⁵ enkephalin of human origin.

PRODUCT

Each vial contains 200 μ g IgG₁ kappa light chain in 1.0 ml of PBS with < 0.1% sodium azide and 0.1% gelatin.

Met/Leu-enkephalin (NOC1/35) is available conjugated to agarose (sc-47705 AC), 500 μ g/0.25 ml agarose in 1 ml, for IP; to HRP (sc-47705 HRP), 200 μ g/ml, for WB, IHC(P) and ELISA; to either phycoerythrin (sc-47705 PE), fluorescein (sc-47705 FITC), Alexa Fluor[®] 488 (sc-47705 AF488), Alexa Fluor[®] 546 (sc-47705 AF546), Alexa Fluor[®] 594 (sc-47705 AF594) or Alexa Fluor[®] 647 (sc-47705 AF647), 200 μ g/ml, for WB (RGB), IF, IHC(P) and FCM; and to either Alexa Fluor[®] 680 (sc-47705 AF680) or Alexa Fluor[®] 790 (sc-47705 AF790), 200 μ g/ml, for Near-Infrared (NIR) WB, IF and FCM.

Alexa Fluor[®] is a trademark of Molecular Probes, Inc., Oregon, USA

APPLICATIONS

Met/Leu-enkephalin (NOC1/35) is recommended for detection of Proenkephalin A and the processed active peptides Met-enkephalin and Leu-enkephalin of mouse, rat and human origin by Western Blotting (starting dilution 1:100, dilution range 1:100-1:1000), immunoprecipitation [1-2 μ g per 100-500 μ g of total protein (1 ml of cell lysate)], immunofluorescence (starting dilution 1:50, dilution range 1:50-1:500) and immunohistochemistry (including paraffin-embedded sections) (starting dilution 1:50, dilution range 1:50-1:500).


STORAGE

Store at 4° C, ****DO NOT FREEZE****. Stable for one year from the date of shipment. Non-hazardous. No MSDS required.


RESEARCH USE

For research use only, not for use in diagnostic procedures.

DATA



Met/Leu-enkephalin (NOC1/35): sc-47705. Western blot analysis of human recombinant Proenkephalin.



Met/Leu-enkephalin (NOC1/35): sc-47705. Immunoperoxidase staining of formalin fixed, paraffin-embedded rat brain tissue showing Enkephalin positive cells in the Ventral Lateral Striatum. Image kindly provided by Regina Vontell, Adrienne Betz and John Salamone, University of Connecticut (A). Immunoperoxidase staining of formalin fixed, paraffin-embedded human duodenum tissue showing cytoplasmic staining of glandular cells. Kindly provided by The Swedish Human Protein Atlas (HPA) program (B).

SELECT PRODUCT CITATIONS

- Huang, W., et al. 2011. Structures and connections of enkephalin- and γ -aminobutyric acid-immunoreactive profiles in the gustatory region of the nucleus tractus solitarius: a light and electron microscopic study. *Neurosci. Biobehav. Rev.* 32: 53-58.
- Shen, H.Y., et al. 2013. Adenosine A_{2A} receptors in striatal glutamatergic terminals and GABAergic neurons oppositely modulate psychostimulant action and DARPP-32 phosphorylation. *PLoS ONE* 8: e80902.
- Kim, J.Y., et al. 2016. Morphine suppresses lung cancer cell proliferation through the interaction with opioid growth factor receptor: an *in vitro* and human lung tissue study. *Anesth. Analg.* 123: 1429-1436.
- Leroy, F., et al. 2017. Input-timing-dependent plasticity in the hippocampal CA2 region and its potential role in social memory. *Neuron* 95: 1089-1102.
- Bosi, G., et al. 2020. Mucosal hallmarks in the alimentary canal of northern pike *Esox lucius* (Linnaeus). *Animals* 10: 1479.
- Cai, H.Y., et al. 2021. Adjusting vascular permeability, leukocyte infiltration, and microglial cell activation to rescue dopaminergic neurons in rodent models of Parkinson's disease. *NPJ Parkinsons Dis.* 7: 91.
- Leroy, F., et al. 2022. Enkephalin release from VIP interneurons in the hippocampal CA2/3a region mediates heterosynaptic plasticity and social memory. *Mol. Psychiatry* 27: 2879-2900.
- Markopoulou, S., et al. 2022. Effect of bladder injection of onabotulinumtoxinA on the central expression of genes associated with the control of the lower urinary tract: a study in normal rats. *Int. J. Mol. Sci.* 23: 14419.

PROTOCOLS

See our web site at www.scbt.com for detailed protocols and support products.