

TCP-1 β (F39 P7 F11): sc-47717

BACKGROUND

The protein TCP-1 (t complex polypeptide 1) is a subunit of the hetero-oligomeric complex CCT (chaperonin containing TCP-1) present in the eukaryotic cytosol. The CCT of eukaryotic cytosol is composed of eight different subunit species, TCP-1 α , β , γ , δ , ϵ , ζ , η and θ , each encoded by a different gene. Two ζ subunits have been described: TCP-1 ζ (also designated TCP-1 ξ 1) and TCP-1 ζ 2. TCP-1 subunits are proposed to have independent functions in folding its *in vivo* substrates, the Actins and Tubulins. TCP-1 was first identified in the mouse as relevant for tail-less and embryonic lethal phenotypes. Sequences homologous to TCP-1 have been isolated in several other species, and the yeast TCP-1 has been shown to encode a molecular chaperone for Actin and Tubulin. TCP-1 found in mammalian cells and yeast plays an important role in the folding of cytosolic proteins.

REFERENCES

1. Ahnert, V., et al. 1996. Cucumber T-complex protein. Molecular cloning, bacterial expression and characterization within a 22-S cytosolic complex in cotyledons and hypocotyls. *Eur. J. Biochem.* 235: 114-119.
2. Iijima, M., et al. 1998. A *Dictyostelium discoideum* homologue to TCP-1 is essential for growth and development. *Gene* 213: 101-106.
3. Ritco-Vonsovici, M. and Willison, K.R. 2000. Defining the eukaryotic cytosolic chaperonin-binding sites in human Tubulins. *J. Mol. Biol.* 304: 81-98.

CHROMOSOMAL LOCATION

Genetic locus: CCT2 (human) mapping to 12q15; Cct2 (mouse) mapping to 10 D2.

SOURCE

TCP-1 β (F39 P7 F11) is a mouse monoclonal antibody raised against ovalbumin-conjugate synthetic peptide RKRVPDHHPC.

PRODUCT

Each vial contains 200 μ g IgG_{2a} kappa light chain in 1.0 ml of PBS with < 0.1% sodium azide and 0.1% gelatin.

APPLICATIONS

TCP-1 β (F39 P7 F11) is recommended for detection of TCP-1 β of mouse, rat, human and *Xenopus laevis* origin by Western Blotting (starting dilution 1:100, dilution range), immunoprecipitation [1-2 μ g per 100-500 μ g of total protein (1 ml of cell lysate)], immunofluorescence (starting dilution 1:50, dilution range 1:50-1:500) and immunohistochemistry (including paraffin-embedded sections) (starting dilution 1:50, dilution range 1:50-1:500).

Suitable for use as control antibody for TCP-1 β siRNA (h): sc-36622, TCP-1 β siRNA (m): sc-36625, TCP-1 β shRNA Plasmid (h): sc-36622-SH, TCP-1 β shRNA Plasmid (m): sc-36625-SH, TCP-1 β shRNA (h) Lentiviral Particles: sc-36622-V and TCP-1 β shRNA (m) Lentiviral Particles: sc-36625-V.


Molecular Weight of TCP-1 β : 50 kDa.

Positive Controls: Caki-1 cell lysate: sc-2224, HEL 92.1.7 cell lysate: sc-2270 or F9 cell lysate: sc-2245.


RECOMMENDED SUPPORT REAGENTS

To ensure optimal results, the following support reagents are recommended: 1) Western Blotting: use m-IgG κ BP-HRP: sc-516102 or m-IgG κ BP-HRP (Cruz Marker): sc-516102-CM (dilution range: 1:1000-1:10000), Cruz Marker™ Molecular Weight Standards: sc-2035, UltraCruz® Blocking Reagent: sc-516214 and Western Blotting Luminol Reagent: sc-2048. 2) Immunoprecipitation: use Protein A/G PLUS-Agarose: sc-2003 (0.5 ml agarose/2.0 ml). 3) Immunofluorescence: use m-IgG κ BP-FITC: sc-516140 or m-IgG κ BP-PE: sc-516141 (dilution range: 1:50-1:200) with UltraCruz® Mounting Medium: sc-24941 or UltraCruz® Hard-set Mounting Medium: sc-359850. 4) Immunohistochemistry: use m-IgG κ BP-HRP: sc-516102 with DAB, 50X: sc-24982 and Immunohistomount: sc-45086, or Organo/Limonene Mount: sc-45087.

DATA



TCP-1 β (F39 P7 F11): sc-47717. Western blot analysis of TCP-1 β expression in HEL 92.1.7 (A), Caki-1 (B), ZR-75-1 (C), F9 (D) and NIH/3T3 (E) whole cell lysates. Detection reagent used: m-IgG_{2a} BP-HRP: sc-542731.



TCP-1 β (F39 P7 F11): sc-47717. Immunofluorescence staining of methanol-fixed HeLa cells showing cytoplasmic localization.

SELECT PRODUCT CITATIONS

1. Song, H. and Sokolov, M. 2009. Analysis of protein expression and compartmentalization in retinal neurons using serial tangential sectioning of the retina. *J. Proteome Res.* 8: 346-351.
2. Tashiro, E., et al. 2013. Prefoldin protects neuronal cells from polyglutamine toxicity by preventing aggregation formation. *J. Biol. Chem.* 288: 19958-19972.
3. Gao, X., et al. 2013. Splice isoforms of phosducin-like protein control the expression of heterotrimeric G proteins. *J. Biol. Chem.* 288: 25760-25768.

STORAGE

Store at 4° C, **DO NOT FREEZE**. Stable for one year from the date of shipment. Non-hazardous. No MSDS required.

RESEARCH USE

For research use only, not for use in diagnostic procedures.

PROTOCOLS

See our web site at www.scbt.com for detailed protocols and support products.