

CTMP (C-18): sc-47857

BACKGROUND

The Akt family of protein kinases (also designated PKB) play a role in Insulin signaling, cellular survival and transformation. Genetic alterations resulting in an aberrant activation of the phosphoinositol-3-kinase (PI3K)/Akt signaling pathway frequently occur in glioblastomas. Such factors include phosphatase and tensin homolog (PTEN) mutation, epidermal growth factor receptor (EGFR) amplification and rearrangement, and carboxy-terminal modulator protein (CTMP) hypermethylation. CTMP binds to the C-terminal regulatory domain of Akt and acts as a negative regulator. CTMP inhibits phosphorylation of Akt on serine 473 and threonine 308, thereby reducing its activity. Decreased expression of CTMP by hypermethylation of its promoter has been linked to the pathogenesis of glioblastomas.

REFERENCES

- Maira, S.M., Galetic, I., Brazil, D.P., Kaech, S., Ingley, E., Thelen, M. and Hemmings, B.A. 2001. Carboxyl-terminal modulator protein (CTMP), a negative regulator of PKB/Akt and v-Akt at the plasma membrane. *Science* 294: 374-380.
- Knobbe, C.B., Reifemberger, J., Blaschke, B. and Reifemberger, G. 2004. Hypermethylation and transcriptional downregulation of the carboxyl-terminal modulator protein gene in glioblastomas. *J. Natl. Cancer Inst.* 96: 483-486.
- Knobbe, C.B., Trampe-Kieslich, A. and Reifemberger, G. 2005. Genetic alteration and expression of the phosphoinositol-3-kinase/Akt pathway genes PIK3CA and PIKE in human glioblastomas. *Neuropathol. Appl. Neurobiol.* 31: 486-490.
- Chae, K.S., Martin-Caraballo, M. anderson, M. and Dryer, S.E. 2005. Akt activation is necessary for growth factor-induced trafficking of functional K(Ca) channels in developing parasympathetic neurons. *J. Neurophysiol.* 93: 1174-1182.
- Williams, D.L., Ozment-Skelton, T. and Li, C. 2006. Modulation of the phosphoinositide 3-kinase signaling pathway alters host response to sepsis, inflammation, and ischemia/reperfusion injury. *Shock* 25: 432-439.
- Martelli, A.M., Faenza, I., Billi, A.M., Manzoli, L., Evangelisti, C., Falà, F. and Cocco, L. 2006. Intracellular 3'-phosphoinositide metabolism and Akt signaling: new mechanisms for tumorigenesis and protection against apoptosis? *Cell. Signal.* 18: 1101-1107.

CHROMOSOMAL LOCATION

Genetic locus: THEM4 (human) mapping to 1q21.3.

SOURCE

CTMP (C-18) is an affinity purified goat polyclonal antibody raised against a peptide mapping at the C-terminus of CTMP of human origin.

STORAGE

Store at 4° C, ****DO NOT FREEZE****. Stable for one year from the date of shipment. Non-hazardous. No MSDS required.

PRODUCT

Each vial contains 200 µg IgG in 1.0 ml of PBS with < 0.1% sodium azide and 0.1% gelatin.

Blocking peptide available for competition studies, sc-47857 P, (100 µg peptide in 0.5 ml PBS containing < 0.1% sodium azide and 0.2% BSA).

APPLICATIONS

CTMP (C-18) is recommended for detection of CTMP isoform a of human origin by Western Blotting (starting dilution 1:200, dilution range 1:100-1:1000), immunofluorescence (starting dilution 1:50, dilution range 1:50-1:500) and solid phase ELISA (starting dilution 1:30, dilution range 1:30-1:3000).

CTMP (C-18) is also recommended for detection of CTMP isoform a in additional species, including bovine.

Suitable for use as control antibody for CTMP siRNA (h): sc-60469, CTMP shRNA Plasmid (h): sc-60469-SH and CTMP shRNA (h) Lentiviral Particles: sc-60469-V.

Molecular Weight of CTMP: 26 kDa.

RECOMMENDED SECONDARY REAGENTS

To ensure optimal results, the following support (secondary) reagents are recommended: 1) Western Blotting: use donkey anti-goat IgG-HRP: sc-2020 (dilution range: 1:2000-1:100,000) or Cruz Marker™ compatible donkey anti-goat IgG-HRP: sc-2033 (dilution range: 1:2000-1:5000), Cruz Marker™ Molecular Weight Standards: sc-2035, TBS Blotto A Blocking Reagent: sc-2333 and Western Blotting Luminol Reagent: sc-2048. 2) Immunofluorescence: use donkey anti-goat IgG-FITC: sc-2024 (dilution range: 1:100-1:400) or donkey anti-goat IgG-TR: sc-2783 (dilution range: 1:100-1:400) with UltraCruz™ Mounting Medium: sc-24941.

RESEARCH USE

For research use only, not for use in diagnostic procedures.

PROTOCOLS

See our web site at www.scbt.com or our catalog for detailed protocols and support products.