

ELKS (N-20): sc-47877

BACKGROUND

The glutamine, leucine, lysine and serine-rich protein ELKS, also designated Rab6-interacting protein 2 and CAZ-associated structural protein (CAST) or the acronym ERC, is a member of a family of RIM-binding proteins. RIMs are presynaptic active zone proteins that regulate neurotransmitter release. This class of protein functions by recruiting I κ B α to the I κ B kinase (IKK) complex, and thus serves a regulatory function for IKK activation. Five isoforms of ELKS (α , β , γ , δ and ϵ) exist and have multiple coding region differences and distinct C-termini. Only brain-specific ELKS bind to RIMs, but both ubiquitous and brain-specific ELKS bind to Rab6, a GTP-binding protein involved in membrane traffic at the Golgi complex. Fusion of ELKS to RET due to translocation t(10;12)(q11;p13) results in a papillary thyroid carcinoma.

CHROMOSOMAL LOCATION

Genetic locus: ERC1 (human) mapping to 12p13.33; Erc1 (mouse) mapping to 6 F1.

SOURCE

ELKS (N-20) is an affinity purified goat polyclonal antibody raised against a peptide mapping near the N-terminus of ELKS of human origin.

PRODUCT

Each vial contains 200 μ g IgG in 1.0 ml of PBS with < 0.1% sodium azide and 0.1% gelatin. Also available as TransCruz reagent for Gel Supershift and ChIP applications, sc-47877 X, 200 μ g/0.1 ml.

Blocking peptide available for competition studies, sc-47877 P, (100 μ g peptide in 0.5 ml PBS containing < 0.1% sodium azide and 0.2% BSA).

RESEARCH USE

For research use only, not for use in diagnostic procedures.

APPLICATIONS

ELKS (N-20) is recommended for detection of ELKS of mouse, rat and human origin by Western Blotting (starting dilution 1:200, dilution range 1:100-1:1000), immunofluorescence (starting dilution 1:50, dilution range 1:50-1:500) and solid phase ELISA (starting dilution 1:30, dilution range 1:30-1:3000).

ELKS (N-20) is also recommended for detection of ELKS in additional species, including equine, canine, bovine, porcine and avian.

Suitable for use as control antibody for ELKS siRNA (h): sc-60572, ELKS siRNA (m): sc-60573, ELKS shRNA Plasmid (h): sc-60572-SH, ELKS shRNA Plasmid (m): sc-60573-SH, ELKS shRNA (h) Lentiviral Particles: sc-60572-V and ELKS shRNA (m) Lentiviral Particles: sc-60573-V.

ELKS (N-20) X TransCruz antibody is recommended for Gel Supershift and ChIP applications.


Molecular Weight of ELKS: 94 kDa.

Positive Controls: MIA PaCa-2 cell lysate: sc-2285, SW480 cell lysate: sc-2219 or mouse testis extract: sc-2405.


RECOMMENDED SECONDARY REAGENTS

To ensure optimal results, the following support (secondary) reagents are recommended: 1) Western Blotting: use donkey anti-goat IgG-HRP: sc-2020 (dilution range: 1:2000-1:100,000) or Cruz Marker[™] compatible donkey anti-goat IgG-HRP: sc-2033 (dilution range: 1:2000-1:5000), Cruz Marker[™] Molecular Weight Standards: sc-2035, TBS Blotto A Blocking Reagent: c-2333 and Western Blotting Luminol Reagent: sc-2048. 2) Immunofluorescence: use donkey anti-goat IgG-FITC: sc-2024 (dilution range: 1:100-1:400) or donkey anti-goat IgG-TR: sc-2783 (dilution range: 1:100-1:400) with UltraCruz[™] Mounting Medium: sc-24941.

DATA



ELKS (N-20): sc-47877. Western blot analysis of ELKS expression in MIA PaCa-2 whole cell lysate.



ELKS (N-20): sc-47877. Western blot analysis of ELKS expression in 293T (A) and SW480 (B) whole cell lysates.

STORAGE

Store at 4° C, ****DO NOT FREEZE****. Stable for one year from the date of shipment. Non-hazardous. No MSDS required.

PROTOCOLS

See our web site at www.scbt.com or our catalog for detailed protocols and support products.



Try **ELKS (G-10): sc-515041** or **ELKS (E-1): sc-365715**, our highly recommended monoclonal alternatives to ELKS (N-20).