

CSN6 (C-17): sc-47965

BACKGROUND

The COP9 signalosome (CSN) complex is involved in several different developmental and cellular processes. The complex is made up of several widely expressed proteins: CSN1 (COPS1), CSN2 (COPS2), CSN3 (COPS3), CSN4 (COPS4), CSN5 (COPS5), CSN6 (COP6), CSN7a (COPS7, COPS7a) or CSN7b (COP7b) and CSN8 (COP8). The CSN complex acts as a regulator for the ubiquitin conjugation pathway by mediating the deneddylation of the SCF-type E3 ligase complexes, which leads to a decrease in ubiquitin ligase activity of SCF-complexes. It is also involved in the phosphorylation of p53, c-Jun, IκBα and IRF-8, as well as CSN-dependent phosphorylation of p53. c-Jun protects and promotes degradation by the Ubl system.

REFERENCES

1. Seeger, M., et al. 1998. A novel protein complex involved in signal transduction possessing similarities to 26S Proteasome subunits. *FASEB J.* 12: 469-478.
2. Bech-Otschir, D., et al. 2001. COP9 signalosome-specific phosphorylation targets p53 to degradation by the ubiquitin system. *EMBO J.* 20: 1630-1639.

CHROMOSOMAL LOCATION

Genetic locus: COPS6 (human) mapping to 7q22.1; Cops6 (mouse) mapping to 5 G2.

SOURCE

CSN6 (C-17) is an affinity purified goat polyclonal antibody raised against a peptide mapping near the C-terminus of CSN6 of human origin.

PRODUCT

Each vial contains 200 µg IgG in 1.0 ml of PBS with < 0.1% sodium azide and 0.1% gelatin.

Blocking peptide available for competition studies, sc-47965 P, (100 µg peptide in 0.5 ml PBS containing < 0.1% sodium azide and 0.2% BSA).

APPLICATIONS

CSN6 (C-17) is recommended for detection of CSN6 of mouse, rat and human origin by Western Blotting (starting dilution 1:200, dilution range 1:100-1:1000), immunoprecipitation [1-2 µg per 100-500 µg of total protein (1 ml of cell lysate)], immunofluorescence (starting dilution 1:50, dilution range 1:50-1:500) and solid phase ELISA (starting dilution 1:30, dilution range 1:30-1:3000).

CSN6 (C-17) is also recommended for detection of CSN6 in additional species, including equine, canine, bovine and porcine.

Suitable for use as control antibody for CSN6 siRNA (h): sc-60461, CSN6 siRNA (m): sc-60462, CSN6 shRNA Plasmid (h): sc-60461-SH, CSN6 shRNA Plasmid (m): sc-60462-SH, CSN6 shRNA (h) Lentiviral Particles: sc-60461-V and CSN6 shRNA (m) Lentiviral Particles: sc-60462-V.


Molecular Weight of CSN6: 34 kDa.

Positive Controls: HeLa whole cell lysate: sc-2200, SW480 cell lysate: sc-2219 or rat brain extract: sc-2392


RECOMMENDED SECONDARY REAGENTS

To ensure optimal results, the following support (secondary) reagents are recommended: 1) Western Blotting: use donkey anti-goat IgG-HRP: sc-2020 (dilution range: 1:2000-1:100,000) or Cruz Marker™ compatible donkey anti-goat IgG-HRP: sc-2033 (dilution range: 1:2000-1:5000), Cruz Marker™ Molecular Weight Standards: sc-2035, TBS Blotto A Blocking Reagent: sc-2333 and Western Blotting Luminol Reagent: sc-2048. 2) Immunoprecipitation: use Protein A/G PLUS-Agarose: sc-2003 (0.5 ml agarose/2.0 ml). 3) Immunofluorescence: use donkey anti-goat IgG-FITC: sc-2024 (dilution range: 1:100-1:400) or donkey anti-goat IgG-TR: sc-2783 (dilution range: 1:100-1:400) with UltraCruz™ Mounting Medium: sc-24941.

DATA



CSN6 (C-17): sc-47965. Western blot analysis of CSN6 expression in HeLa (A), SW480 (B), 3T3-L1 (C) and A-375 (D) whole cell lysates and mouse brain tissue extract (E).



CSN6 (C-17): sc-47965. Western blot analysis of CSN6 expression in rat brain tissue extract.

STORAGE

Store at 4° C, **DO NOT FREEZE**. Stable for one year from the date of shipment. Non-hazardous. No MSDS required.

RESEARCH USE

For research use only, not for use in diagnostic procedures.

PROTOCOLS

See our web site at www.scbt.com or our catalog for detailed protocols and support products.

MONOS
Satisfaction
Guaranteed

Try **CSN6 (B-1): sc-393023** or **CSN6 (H-3): sc-137122**, our highly recommended monoclonal alternatives to CSN6 (C-17).