

GP73 (C-15): sc-48011

BACKGROUND

GP73 (also known as Golgi phosphoprotein 2, GOLPH 2 or Golgi membrane protein) is a widely expressed, epithelial-specific, type II transmembrane protein which resides in the Golgi apparatus, where it is responsible for the posttranslational modification of proteins produced in the rough ER while assisting in the transport of proteins through the Golgi. The human GP73 gene has been mapped within a BAC and localized to chromosome 9q21.33. GP73 levels rise in those who have been diagnosed with acute and chronic liver diseases.

REFERENCES

1. Kladney, R.D., et al. 2000. GP73, a novel Golgi-localized protein upregulated by viral infection. *Gene* 249: 53-65.
2. Kladney, R.D., et al. 2002. Expression of GP73, a resident Golgi membrane protein, in viral and nonviral liver disease. *Hepatology* 35: 1431-1440.
3. Kladney, R.D., et al. 2002. Upregulation of the Golgi protein GP73 by adenovirus infection requires the E1A CtBP interaction domain. *Virology* 301: 236-246.
4. Iftikhar, R., et al. 2004. Disease- and cell-specific expression of GP73 in human liver disease. *Am. J. Gastroenterol.* 99: 1087-1095.

CHROMOSOMAL LOCATION

Genetic locus: Golp2 (mouse) mapping to 13 B2.

SOURCE

GP73 (C-15) is an affinity purified goat polyclonal antibody raised against a peptide mapping at the C-terminus of GP73 of mouse origin.

PRODUCT

Each vial contains 200 µg IgG in 1.0 ml of PBS with < 0.1% sodium azide and 0.1% gelatin.

Blocking peptide available for competition studies, sc-48011 P, (100 µg peptide in 0.5 ml PBS containing < 0.1% sodium azide and 0.2% BSA).

APPLICATIONS

GP73 (C-15) is recommended for detection of GP73 of mouse origin by Western Blotting (starting dilution 1:200, dilution range 1:100-1:1000), immunoprecipitation [1-2 µg per 100-500 µg of total protein (1 ml of cell lysate)], immunofluorescence (starting dilution 1:50, dilution range 1:50-1:500), immunohistochemistry (including paraffin-embedded sections) (starting dilution 1:50, dilution range 1:50-1:500) and solid phase ELISA (starting dilution 1:30, dilution range 1:30-1:3000).

Suitable for use as control antibody for GP73 siRNA (m): sc-60712, GP73 shRNA Plasmid (m): sc-60712-SH and GP73 shRNA (m) Lentiviral Particles: sc-60712-V.


Molecular Weight of GP73: 73 kDa.

Positive Controls: c4 whole cell lysate: sc-364186 or mouse prostate extract: sc-364249.


RECOMMENDED SECONDARY REAGENTS

To ensure optimal results, the following support (secondary) reagents are recommended: 1) Western Blotting: use donkey anti-goat IgG-HRP: sc-2020 (dilution range: 1:2000-1:100,000) or Cruz Marker™ compatible donkey anti-goat IgG-HRP: sc-2033 (dilution range: 1:2000-1:5000), Cruz Marker™ Molecular Weight Standards: sc-2035, TBS Blotto A Blocking Reagent: sc-2333 and Western Blotting Luminol Reagent: sc-2048. 2) Immunoprecipitation: use Protein A/G PLUS-Agarose: sc-2003 (0.5 ml agarose/2.0 ml). 3) Immunofluorescence: use donkey anti-goat IgG-FITC: sc-2024 (dilution range: 1:100-1:400) or donkey anti-goat IgG-TR: sc-2783 (dilution range: 1:100-1:400) with UltraCruz™ Mounting Medium: sc-24941. 4) Immunohistochemistry: use ImmunoCruz™: sc-2053 or ABC: sc-2023 goat IgG Staining Systems.

DATA



GP73 (C-15): sc-48011. Western blot analysis of GP73 expression in c4 whole cell lysate.



GP73 (C-15): sc-48011. Immunoperoxidase staining of formalin fixed, paraffin-embedded human prostate tissue showing cytoplasmic staining of glandular cells.

STORAGE

Store at 4° C, ****DO NOT FREEZE****. Stable for one year from the date of shipment. Non-hazardous. No MSDS required.

RESEARCH USE

For research use only, not for use in diagnostic procedures.

PROTOCOLS

See our web site at www.scbt.com or our catalog for detailed protocols and support products.



Try **GP73 (E-7): sc-393935** or **GP73 (G-2): sc-398230**, our highly recommended monoclonal alternatives to GP73 (C-15). Also, for AC, HRP, FITC, PE, Alexa Fluor® 488 and Alexa Fluor® 647 conjugates, see **GP73 (E-7): sc-393935**.