

MELK (N-19): sc-48037



The Power to Question

BACKGROUND

Maternal embryonic leucine zipper kinase (KIAA0175, HPK38) or MELK, a new member of the Snf1/AMPK family of kinases, encodes a protein with a kinase catalytic domain and a leucine zipper motif consisting of a periodic repetition of leucine residues at every seventh residue located within the N-terminal catalytic domain. This motif has been observed in myriad DNA-binding proteins and is presumed to be involved in protein-DNA interactions, and potentially protein-protein interactions. Research predicts that the gene product of MELK plays a role in the signal transduction events in the egg and early embryo. Mouse and human MELK proteins share 95% sequence identity in the kinase domain and northern blot analysis in mouse indicates that MELK expression is restricted to spermatogonia in the testis and to oocytes in the ovary.

REFERENCES

- Nagase, T., Seki, N., Ishikawa, K., Tanaka, A. and Nomura, N. 1996. Prediction of the coding sequences of unidentified human genes. V. The coding sequences of 40 new genes (KIAA0161-KIAA0200) deduced by analysis of cDNA clones from human cell line KG-1. *DNA Res.* 3: 17-24.
- Heyer, B.S., Warsowe, J., Solter, D., Knowles, B.B. and Ackerman, S.L. 1997. New member of the Snf1/AMPK kinase family, Melk, is expressed in the mouse egg and preimplantation embryo. *Mol. Reprod. Dev.* 47: 148-156.
- Seong, H.A., Gil, M., Kim, K.T., Kim, S.J. and Ha, H. 2002. Phosphorylation of a novel zinc-finger-like protein, ZPR9, by murine protein serine/threonine kinase 38 (MPK38). *Biochem. J.* 361: 597-604.
- Online Mendelian Inheritance in Man, OMIM™. 2002. Johns Hopkins University, Baltimore, MD. MIM Number: 607025. World Wide Web URL: <http://www.ncbi.nlm.nih.gov/omim/>
- Vulsteke, V., Beullens, M., Boudrez, A., Keppens, S., Van Eynde, A., Rider, M.H., Stalmans, W. and Bollen, M. 2004. Inhibition of spliceosome assembly by the cell cycle-regulated protein kinase MELK and involvement of splicing factor NIPP1. *J. Biol. Chem.* 279: 8642-8647.
- Beullens, M., Vancauwenbergh, S., Morrice, N., Derua, R., Ceulemans, H., Waelkens, E. and Bollen, M. 2005. Substrate specificity and activity regulation of protein kinase MELK. *J. Biol. Chem.* 280: 40003-40011.
- Nakano, I., Paucar, A.A., Bajpai, R., Dougherty, J.D., Zewail, A., Kelly, T.K., Kim, K.J., Ou, J., Groszer, M., Imura, T., Freije, W.A., Nelson, S.F., Sofroniew, M.V., Wu, H., Liu, X., Tersikh, A.V., Geschwind, D.H. and Kornblum, H.I. 2005. Maternal embryonic leucine zipper kinase (MELK) regulates multipotent neural progenitor proliferation. *J. Cell Biol.* 170: 413-427.
- Badouel, C., Körner, R., Frank-Vaillant, M., Couturier, A., Nigg, EA. and Tassan, J.P. 2006. M-phase MELK activity is regulated by MPF and MAPK. *Cell Cycle* 5: 883-889.
- Cordes, S., Frank, C.A. and Garriga, G. 2006. The *C. elegans* MELK ortholog PIG-1 cell fate in asymmetric neuroblast divisions. *Dev.* 133: 2747-2756.

CHROMOSOMAL LOCATION

Genetic locus: MELK (human) mapping to 9p13.2; Melk (mouse) mapping to 4 B1.

SOURCE

MELK (N-19) is an affinity purified goat polyclonal antibody raised against a peptide mapping at the N-terminus of MELK of human origin.

PRODUCT

Each vial contains 200 µg IgG in 1.0 ml of PBS with < 0.1% sodium azide and 0.1% gelatin.

Blocking peptide available for competition studies, sc-48037 P, (100 µg peptide in 0.5 ml PBS containing < 0.1% sodium azide and 0.2% BSA).

APPLICATIONS

MELK (N-19) is recommended for detection of MELK of mouse, rat and human origin by Western Blotting (starting dilution 1:200, dilution range 1:100-1:1000), immunofluorescence (starting dilution 1:50, dilution range 1:50-1:500) and solid phase ELISA (starting dilution 1:30, dilution range 1:30-1:3000).

Suitable for use as control antibody for MELK siRNA (h): sc-61016 and MELK siRNA (m): sc-61017.

Molecular Weight of MELK: 73 kDa.

RECOMMENDED SECONDARY REAGENTS

To ensure optimal results, the following support (secondary) reagents are recommended: 1) Western Blotting: use donkey anti-goat IgG-HRP: sc-2020 (dilution range: 1:2000-1:100,000) or Cruz Marker™ compatible donkey anti-goat IgG-HRP: sc-2033 (dilution range: 1:2000-1:5000), Cruz Marker™ Molecular Weight Standards: sc-2035, TBS Blotto A Blocking Reagent: sc-2333 and Western Blotting Luminol Reagent: sc-2048. 2) Immunofluorescence: use donkey anti-goat IgG-FITC: sc-2024 (dilution range: 1:100-1:400) or donkey anti-goat IgG-TR: sc-2783 (dilution range: 1:100-1:400) with UltraCruz™ Mounting Medium: sc-24941.

STORAGE

Store at 4° C, **DO NOT FREEZE**. Stable for one year from the date of shipment. Non-hazardous. No MSDS required.

RESEARCH USE

For research use only, not for use in diagnostic procedures.

PROTOCOLS

See our web site at www.scbt.com or our catalog for detailed protocols and support products.