

NIS (S-16): sc-48058

BACKGROUND

The sodium/iodide symporter (NIS) is an integral plasma membrane glycoprotein that mediates active iodide transport in the thyroid and other tissues, including salivary glands, gastric mucosa and lactating mammary gland. In the lactating mammary gland, NIS transports iodide into the milk, thereby allowing the nursing newborn to use the iodide for thyroid hormone biosynthesis. NIS is expressed in some breast cancers, but exhibits decreased expression in the majority of thyroid cancers, most likely due to alterations in the binding activity of AP2 and Sp1 transcription factors to the NIS promoter. NIS is a prerequisite for radioiodide treatment of thyroid cancer and a promising diagnostic and therapeutic tool for breast cancer.

REFERENCES

1. Paulini, K. and Mohr, W. 1975. Hormone-dependent polyploidy in the glandula orbitalis externa and glandula infraorbitalis of animals of different age. *Beitr. Pathol.* 156: 65-74.
2. Boismare, F., Le Poncin, M., Lefrançois, J., Hacpille, L. and Marchand, J.C. 1977. The treatment, by Imipramine, of the hemodynamic, functional and biochemical consequences, of an experimental cranio-cervical trauma in rats. *C. R. Seances Soc. Biol. Fil.* 170: 1110-1117.
3. Partona, F., Djakaria, Oemijati, S., Joesoef, A., Clarke, M.D., Cole, W.C., Lien, J.C. and Cross, J.H. 1978. Filariasis in West Kalimantan (Borneo), Indonesia. *Southeast Asian J. Trop. Med. Public Health* 8: 459-463.
4. Osteen, K.G. and Mills, T.M. 1980. Serum LH and FSH levels in the pregnant rabbit. *Proc. Soc. Exp. Biol. Med.* 162: 454-457.
5. Kogai, T., Kanamoto, Y., Li, A.I., Che, L.H., Ohashi, E., Taki, K., Chandraratna, R.A., Saito, T. and Brent, G.A. 2005. Differential regulation of sodium/iodide symporter gene expression by nuclear receptor ligands in MCF-7 breast cancer cells. *Endocrinology* 146: 3059-3069.
6. Miyagawa, M., Anton, M., Wagner, B., Haubner, R., Souvatzoglou, M., Gansbacher, B., Schwaiger, M. and Bengel, F.M. 2005. Non-invasive imaging of cardiac transgene expression with PET: comparison of the human sodium/iodide symporter gene and HSV1-tk as the reporter gene. *Eur. J. Nucl. Med. Mol. Imaging* 32: 1108-1114.
7. Schmitz, G., Füzesi, L., Struck, J., Siefker, U., Hamann, A., Sahlmann, C.O., Hufner, M. and Meller, J. 2005. Expression of the sodium iodide symporter in differentiated thyroid cancer: clinical evidence. *Nuklearmedizin* 44: 86-93.

CHROMOSOMAL LOCATION

Genetic locus: Slc5a5 (mouse) mapping to 8 B3.3.

SOURCE

NIS (S-16) is an affinity purified goat polyclonal antibody raised against a peptide mapping within an extracellular domain of NIS of mouse origin.

STORAGE

Store at 4° C, ****DO NOT FREEZE****. Stable for one year from the date of shipment. Non-hazardous. No MSDS required.

PRODUCT

Each vial contains 200 µg IgG in 1.0 ml of PBS with < 0.1% sodium azide and 0.1% gelatin.

Blocking peptide available for competition studies, ready P, (100 µg peptide in 0.5 ml PBS containing < 0.1% sodium azide and 0.2% BSA).

APPLICATIONS

NIS (S-16) is recommended for detection of NIS of mouse and, to a lesser extent, rat origin by Western Blotting (starting dilution 1:200, dilution range 1:100-1:1000), immunofluorescence (starting dilution 1:50, dilution range 1:50-1:500) and solid phase ELISA (starting dilution 1:30, dilution range 1:30-1:3000).

Suitable for use as control antibody for NIS siRNA (m): sc-61200, NIS shRNA Plasmid (m): sc-61200-SH and NIS shRNA (m) Lentiviral Particles: sc-61200-V.

Molecular Weight of non-glycosylated NIS: 50 kDa.

Molecular Weight of glycosylated NIS: 87-110 kDa.

RECOMMENDED SECONDARY REAGENTS

To ensure optimal results, the following support (secondary) reagents are recommended: 1) Western Blotting: use donkey anti-goat IgG-HRP: sc-2020 (dilution range: 1:2000-1:100,000) or Cruz Marker™ compatible donkey anti-goat IgG-HRP: sc-2033 (dilution range: 1:2000-1:5000), Cruz Marker™ Molecular Weight Standards: sc-2035, TBS Blotto A Blocking Reagent: sc-2333 and Western Blotting Luminol Reagent: sc-2048. 2) Immunofluorescence: use donkey anti-goat IgG-FITC: sc-2024 (dilution range: 1:100-1:400) or donkey anti-goat IgG-TR: sc-2783 (dilution range: 1:100-1:400) with UltraCruz™ Mounting Medium: sc-24941.

SELECT PRODUCT CITATIONS

1. Gaertner, F.C., Rohde, F., Mueller, J., Blechert, B., Janssen, K.P. and Essler, M. 2009. Endogenous expression of the sodium iodide symporter mediates uptake of iodide in murine models of colorectal carcinoma. *Int. J. Cancer* 125: 2783-2791.

RESEARCH USE

For research use only, not for use in diagnostic procedures.

PROTOCOLS

See our web site at www.scbt.com or our catalog for detailed protocols and support products.



Try **NIS (G-5): sc-514487**, our highly recommended monoclonal alternative to NIS (S-16).