

# TEM7 (N-19): sc-48103

## BACKGROUND

Tumor endothelial markers (TEMs) are abundantly expressed in the blood vessels of human solid tumors during angiogenesis and neoangiogenesis. These include TEM1 (endosialin), TEM5 (G protein-coupled receptor 124) and TEM7 (plexin domain containing 1). TEMs are associated with the cell surface membrane at low levels in normal human and mouse tissues. TEM5 is a seven-pass transmembrane receptor, whereas TEM1, TEM7 and TEM8 span the membrane once. TEM5 expression is elevated during tumor angiogenesis and neoangiogenesis. TEM7 is highly expressed in tumor endothelium and neurons. Therefore, TEM5 and TEM7 may be suitable targets for the development of antiangiogenic therapies.

## REFERENCES

- Carson-Walter, E.B., Watkins, D.N., Nanda, A., Vogelstein, B., Kinzler, K.W. and St. Croix, B. 2001. Cell surface tumor endothelial markers are conserved in mice and humans. *Cancer Res.* 61: 6649-6655.
- Nanda, A., Buckhaults, P., Seaman, S., Agrawal, N., Boutin, P., Shankara, S., Nacht, M., Teicher, B., Stampfl, J., Singh, S., Vogelstein, B., Kinzler, KW. and St. Croix, B. 2004. Identification of a binding partner for the endothelial cell surface proteins TEM7 and TEM7R. *Cancer Res.* 64: 8507-8511.
- Yamamoto, Y., Irie, K., Asada, M., Mino, A., Mandai, K. and Takai, Y. 2004. Dir gene to seven-pass transmembrane proteins, tumor endothelial marker 5 (TEM5), and a novel TEM5-like protein. *Oncogene* 23: 3889-3897.
- Wang, X.Q., Sheibani, N. and Watson, J.C. 2005. Modulation of tumor endothelial c cell capillary morphogenesis. *Microvasc. Res.* 70: 189-197.
- Lee, H.K., Kang, D.S., Seo, I.A., Choi, E.J., Park, H.T. and Park, J.I. 2006. Expression of tumor endothelial marker 7 mRNA and protein in the dorsal root ganglion neurons of the rat. *Neurosci. Lett.* 402: 71-75.
- Lee, H.K., Seo, I.A., Park, H.K. and Park, H.T. 2006. Identification of the basement membrane protein nidogen as a candidate ligand for tumor endothelial marker 7 *in vitro* and *in vivo*. *FEBS Lett.* 580: 2253-2257.

## CHROMOSOMAL LOCATION

Genetic locus: PLXDC1 (human) mapping to 17q12; Plxdc1 (mouse) mapping to 11 D.

## SOURCE

TEM7 (N-19) is an affinity purified goat polyclonal antibody raised against a peptide mapping near the N-terminus of TEM7 of human origin.

## PRODUCT

Each vial contains 200 µg IgG in 1.0 ml of PBS with < 0.1% sodium azide and 0.1% gelatin.

Blocking peptide available for competition studies, sc-48103 P, (100 µg peptide in 0.5 ml PBS containing < 0.1% sodium azide and 0.2% BSA).

## RESEARCH USE

For research use only, not for use in diagnostic procedures.

## APPLICATIONS

TEM7 (N-19) is recommended for detection of TEM7 of human and, to a lesser extent, mouse origin by Western Blotting (starting dilution 1:200, dilution range 1:100-1:1000), immunoprecipitation [1-2 µg per 100-500 µg of total protein (1 ml of cell lysate)], immunofluorescence (starting dilution 1:50, dilution range 1:50-1:500) and solid phase ELISA (starting dilution 1:30, dilution range 1:30-1:3000).

TEM7 (N-19) is also recommended for detection of TEM7 in additional species, including equine.

Suitable for use as control antibody for TEM7 siRNA (h): sc-61663, TEM7 siRNA (m): sc-61664, TEM7 shRNA Plasmid (h): sc-61663-SH, TEM7 shRNA Plasmid (m): sc-61664-SH, TEM7 shRNA (h) Lentiviral Particles: sc-61663-V and TEM7 shRNA (m) Lentiviral Particles: sc-61664-V.


Molecular Weight of TEM7: 60 kDa.

Positive Controls: HCT-116 whole cell lysate: sc-364175, U-87 MG cell lysate: sc-2411 or ECV304 cell lysate: sc-2269.

## RECOMMENDED SECONDARY REAGENTS

To ensure optimal results, the following support (secondary) reagents are recommended: 1) Western Blotting: use donkey anti-goat IgG-HRP: sc-2020 (dilution range: 1:2000-1:100,000) or Cruz Marker™ compatible donkey anti-goat IgG-HRP: sc-2033 (dilution range: 1:2000-1:5000), Cruz Marker™ Molecular Weight Standards: sc-2035, TBS Blotto A Blocking Reagent: sc-2333 and Western Blotting Luminol Reagent: sc-2048. 2) Immunoprecipitation: use Protein A/G PLUS-Agarose: sc-2003 (0.5 ml agarose/2.0 ml). 3) Immunofluorescence: use donkey anti-goat IgG-FITC: sc-2024 (dilution range: 1:100-1:400) or donkey anti-goat IgG-TR: sc-2783 (dilution range: 1:100-1:400) with UltraCruz™ Mounting Medium: sc-24941.

## DATA



TEM7 (N-19): sc-48103. Western blot analysis of TEM7 expression in 293T whole cell lysate.

## STORAGE

Store at 4° C, **\*\*DO NOT FREEZE\*\***. Stable for one year from the date of shipment. Non-hazardous. No MSDS required.

## PROTOCOLS

See our web site at [www.scbt.com](http://www.scbt.com) or our catalog for detailed protocols and support products.