

Glut6 (M-16): sc-48171

BACKGROUND

The oxidation of glucose functions as the dominant source of metabolic energy for mammals. The plasma membrane is impermeable to glucose, so the cellular uptake of this important nutrient is achieved by facultative hexose transporters (Gluts). Gluts are integral membrane proteins that transport glucose and related hexoses. Glucose binds to a Glut on one side of the membrane which provokes a conformational change causing it to release glucose to the other side. Members of the Glut family may enhance the metabolic activity of tumor cells. Glut6 is part of the third out of three classes of Gluts. Glut6 is mainly expressed in the brain, spleen and peripheral leukocytes. It appears to be regulated by subcellular redistribution, because it is targeted to intracellular compartments by di-leucine motifs, recycling itself in a dynamine-dependent manner.

REFERENCES

- Bell, G.I., Kayano, T., Buse, J.B., Burant, C.F., Takeda, J., Lin, D., Fukumoto, H. and Seino, S. 1990. Molecular biology of mammalian glucose transporters. *Diabetes Care* 13: 198-208.
- Kayano, T., Burant, C.F., Fukumoto, H., Gould, G.W., Fan, Y.S., Eddy, R.L., Byers, M.G., Shows, T.B., Seino, S. and Bell, G.I. 1990. Human facilitative glucose transporters. Isolation, functional characterization, and gene localization of cDNAs encoding an isoform (Glut5) expressed in small intestine, kidney, muscle, and adipose tissue and an unusual glucose transporter pseudogene-like sequence (Glut6). *J. Biol. Chem.* 265: 13276-13282.
- Lisinski, I., Schürmann, A., Joost, H.G., Cushman, S.W. and Al-Hasani, H. 2001. Targeting of Glut6 (formerly Glut9) and Glut8 in rat adipose cells. *Biochem. J.* 358: 517-522.
- Joost, H.G. and Thorens, B. 2002. The extended Glut-family of sugar/polyol transport facilitators: nomenclature, sequence characteristics, and potential function of its novel members (review). *Mol. Membr. Biol.* 18: 247-256.
- Scheepers, A., Joost, H.G. and Schürmann, A. 2004. The glucose transporter families SGLT and Glut: molecular basis of normal and aberrant function. *JPEN J. Parenter. Enteral Nutr.* 28: 364-371.
- Kono, T., Nishida, M., Nishiki, Y., Seki, Y., Sato, K. and Akiba, Y. 2005. Characterisation of glucose transporter (Glut) gene expression in broiler chickens. *Br. Poult. Sci.* 46: 510-515.
- Macheda, M.L., Rogers, S. and Best, J.D. 2005. Molecular and cellular regulation of glucose transporter (Glut) proteins in cancer. *J. Cell. Physiol.* 202: 654-662.

CHROMOSOMAL LOCATION

Genetic locus: SLC2A6 (human) mapping to 9q34.2; Slc2a6 (mouse) mapping to 2 A3.

SOURCE

Glut6 (M-16) is an affinity purified goat polyclonal antibody raised against a peptide mapping within an internal region of Glut6 of mouse origin.

PRODUCT

Each vial contains 200 µg IgG in 1.0 ml of PBS with < 0.1% sodium azide and 0.1% gelatin.

Blocking peptide available for competition studies, sc-48171 P, (100 µg peptide in 0.5 ml PBS containing < 0.1% sodium azide and 0.2% BSA).

APPLICATIONS

Glut6 (M-16) is recommended for detection of Glut6 of mouse and human origin by Western Blotting (starting dilution 1:200, dilution range 1:100-1:1000), immunofluorescence (starting dilution 1:50, dilution range 1:50-1:500) and solid phase ELISA (starting dilution 1:30, dilution range 1:30-1:3000).

Glut6 (M-16) is also recommended for detection of Glut6 in additional species, including porcine.

Suitable for use as control antibody for Glut6 siRNA (h): sc-60699, Glut6 siRNA (m): sc-60700, Glut6 shRNA Plasmid (h): sc-60699-SH, Glut6 shRNA Plasmid (m): sc-60700-SH, Glut6 shRNA (h) Lentiviral Particles: sc-60699-V and Glut6 shRNA (m) Lentiviral Particles: sc-60700-V.

Molecular Weight of Glut6: 55 kDa.

Positive Controls: T24 cell lysate: sc-2292 or MIA PaCa-2 cell lysate: sc-2285.

RECOMMENDED SECONDARY REAGENTS

To ensure optimal results, the following support (secondary) reagents are recommended: 1) Western Blotting: use donkey anti-goat IgG-HRP: sc-2020 (dilution range: 1:2000-1:100,000) or Cruz Marker™ compatible donkey anti-goat IgG-HRP: sc-2033 (dilution range: 1:2000-1:5000), Cruz Marker™ Molecular Weight Standards: sc-2035, TBS Blotto A Blocking Reagent: sc-2333 and Western Blotting Luminol Reagent: sc-2048. 2) Immunofluorescence: use donkey anti-goat IgG-FITC: sc-2024 (dilution range: 1:100-1:400) or donkey anti-goat IgG-TR: sc-2783 (dilution range: 1:100-1:400) with UltraCruz™ Mounting Medium: sc-24941.

STORAGE

Store at 4° C, **DO NOT FREEZE**. Stable for one year from the date of shipment. Non-hazardous. No MSDS required.

RESEARCH USE

For research use only, not for use in diagnostic procedures.


 MONOS
Satisfation
Guaranteed

Try **Glut6 (B-3): sc-373973**, our highly recommended monoclonal alternative to Glut6 (M-16).