

GPR12 (I-18): sc-48199

BACKGROUND

G protein-coupled receptors (GPRs) are a protein family of transmembrane receptors that transmit an extracellular signal (ligand binding) into an intracellular signal (G protein activation). GPR signaling is an evolutionarily ancient mechanism used by all eukaryotes to sense environmental stimuli and mediate cell-cell communication. All of the receptors have seven membrane-spanning domains and the extracellular parts of the receptor can be glycosylated. These extracellular loops also contain two highly conserved cysteine residues which create disulfide bonds to stabilize the receptor structure. GPR12 is a 334 amino acid peptide that is expressed primarily in brain, particularly in regions where neuronal differentiation takes place. GPR12 is coupled to an inhibitory G protein. It positively influences differentiation and maturation of post-mitotic neurons, and it may promote the growth of neuronal precursor cells.

REFERENCES

1. Two new fluoroquinolones. 1992. *Med. Lett. Drugs Ther.* 34: 58-60.
2. Song, Z.H., Modi, W. and Bonner, T.I. 1996. Molecular cloning and chromosomal localization of human genes encoding three closely related G protein-coupled receptors. *Genomics* 28: 347-349.
3. Online Mendelian Inheritance in Man, OMIM[™]. 2002. Johns Hopkins University, Baltimore, MD. MIM Number: 600752. World Wide Web URL: <http://www.ncbi.nlm.nih.gov/omim/>
4. Ignatov, A., Lintzel, J., Hermans-Borgmeyer, I., Kreienkamp, H.J., Joost, P., Thomsen, S., Methner, A. and Schaller, H.C. 2003. Role of the G protein-coupled receptor GPR12 as high-affinity receptor for sphingosylphosphorylcholine and its expression and function in brain development. *J. Neurosci.* 23: 907-914.
5. Uhlenbrock, K., Huber, J., Ardati, A., Busch, A.E. and Kostenis, E. 2003. Fluid shear stress differentially regulates GPR3, GPR6, and GPR12 expression in human umbilical vein endothelial cells. *Cell Physiol. Biochem.* 13: 75-84.
6. Hinckley, M., Vaccari, S., Horner, K., Chen, R. and Conti, M. 2005. The G protein-coupled receptors GPR3 and GPR12 are involved in cAMP signaling and maintenance of meiotic arrest in rodent oocytes. *Dev. Biol.* 287: 249-261.

CHROMOSOMAL LOCATION

Genetic locus: GPR12 (human) mapping to 13q12.13; Gpr12 (mouse) mapping to 5 G3.

SOURCE

GPR12 (I-18) is an affinity purified goat polyclonal antibody raised against a peptide mapping within a cytoplasmic domain of GPR12 of human origin.

STORAGE

Store at 4° C, ****DO NOT FREEZE****. Stable for one year from the date of shipment. Non-hazardous. No MSDS required.

RESEARCH USE

For research use only, not for use in diagnostic procedures.

PRODUCT

Each vial contains 200 µg IgG in 1.0 ml of PBS with < 0.1% sodium azide and 0.1% gelatin.

Blocking peptide available for competition studies, sc-48199 P, (100 µg peptide in 0.5 ml PBS containing < 0.1% sodium azide and 0.2% BSA).

APPLICATIONS

GPR12 (I-18) is recommended for detection of GPR12 of mouse, rat and human origin by Western Blotting (starting dilution 1:200, dilution range 1:100-1:1000), immunofluorescence (starting dilution 1:50, dilution range 1:50-1:500) and solid phase ELISA (starting dilution 1:30, dilution range 1:30-1:3000).

GPR12 (I-18) is also recommended for detection of GPR12 in additional species, including equine, canine, bovine, porcine and avian.

Suitable for use as control antibody for GPR12 siRNA (h): sc-60735, GPR12 siRNA (m): sc-60736, GPR12 shRNA Plasmid (h): sc-60735-SH, GPR12 shRNA Plasmid (m): sc-60736-SH, GPR12 shRNA (h) Lentiviral Particles: sc-60735-V and GPR12 shRNA (m) Lentiviral Particles: sc-60736-V.

Molecular Weight of GPR12: 37 kDa.

Positive Controls: Y79 cell lysate: sc-2240.

RECOMMENDED SECONDARY REAGENTS

To ensure optimal results, the following support (secondary) reagents are recommended: 1) Western Blotting: use donkey anti-goat IgG-HRP: sc-2020 (dilution range: 1:2000-1:100,000) or Cruz Marker[™] compatible donkey anti-goat IgG-HRP: sc-2033 (dilution range: 1:2000-1:5000), Cruz Marker[™] Molecular Weight Standards: sc-2035, TBS Blotto A Blocking Reagent: sc-2333 and Western Blotting Luminol Reagent: sc-2048. 2) Immunofluorescence: use donkey anti-goat IgG-FITC: sc-2024 (dilution range: 1:100-1:400) or donkey anti-goat IgG-TR: sc-2783 (dilution range: 1:100-1:400) with UltraCruz[™] Mounting Medium: sc-24941.

PROTOCOLS

See our web site at www.scbt.com or our catalog for detailed protocols and support products.