

MAGOH (C-12): sc-48237

BACKGROUND

MAGOH, the human homolog of *Drosophila mago nashi*, is required for embryo development. MAGOH is ubiquitously expressed in adult tissues. It has an unusual structure consisting of an extremely flat, six-stranded anti-parallel β sheet packed next to two helices. MAGOH interacts with Y14 to form a complex that plays an crucial role in postslicing processing (including nuclear export and cytoplasmic localization of the mRNA) and in the nonsense-mediated mRNA decay (NMD) surveillance process. The MAGOH-Y14 complex remains persistently associated in the same position on the mRNA after its export to the cytoplasm and requires translation of the mRNA for removal. This complex may illustrate the pre-mRNA splicing machinery's mechanism for forming a stable exon-exon junction complex-mRNA at splice junctions.

REFERENCES

- Zhao, X.F., Colaizzo-Anas, T., Nowak, N.J., Shows, T.B., Elliott, R.W. and Aplan, P.D. 1998. The mammalian homolog of mago nashi encodes a serum-inducible protein. *Genomics* 47: 319-322.
- Zhao, X.F., Nowak, N.J., Shows, T.B. and Aplan, P.D. 2000. MAGOH interacts with a novel RNA-binding protein. *Genomics* 63: 145-148.
- Kataoka, N., Diem, M.D., Kim, V.N., Yong, J. and Dreyfuss, G. 2001. MAGOH, a human homolog of *Drosophila mago nashi* protein, is a component of the splicing-dependent exon-exon junction complex. *EMBO J.* 20: 6424-6433.
- Lau, C.K., Diem, M.D., Dreyfuss, G. and Van Duyne, G.D. 2003. Structure of the MAGOH-Y14 core of the exon junction complex. *Curr. Biol.* 13: 933-941.
- Degot, S., Le Hir, H., Alpy, F., Kedinger, V., Stoll, I., Wendling, C., Seraphin, B., Rio, M.C. and Tomasetto, C. 2004. Association of the breast cancer protein MLN51 with the exon junction complex via its speckle localizer and RNA binding module. *J. Biol. Chem.* 279: 33702-33715.
- Nott, A., Le Hir, H. and Moore, M.J. 2004. Splicing enhances translation in mammalian cells: an additional function of the exon junction complex. *Genes Dev.* 18: 210-222.
- Palacios, I.M., Gatfield, D., St Johnston, D. and Izaurralde, E. 2004. An eIF4AIII-containing complex required for mRNA localization and nonsense-mediated mRNA decay. *Nature* 427: 753-757.
- Wilkinson, M.F. 2005. A new function for nonsense-mediated mRNA-decay factors. *Trends Genet.* 21: 143-148.

CHROMOSOMAL LOCATION

Genetic locus: MAGOH (human) mapping to 1p32.3; Magoh (mouse) mapping to 4 C7, Magohb (mouse) mapping to 6 C7.

SOURCE

MAGOH (C-12) is an affinity purified goat polyclonal antibody raised against a peptide mapping near the C-terminus of MAGOH of human origin.

RESEARCH USE

For research use only, not for use in diagnostic procedures.

PRODUCT

Each vial contains 200 μ g IgG in 1.0 ml of PBS with < 0.1% sodium azide and 0.1% gelatin. Also available as TransCruz reagent for Gel Supershift and ChIP applications, sc-48237 X, 200 μ g/0.1 ml.

Blocking peptide available for competition studies, sc-48237 P, (100 μ g peptide in 0.5 ml PBS containing < 0.1% sodium azide and 0.2% BSA).

APPLICATIONS

MAGOH (C-12) is recommended for detection of MAGOH of human, mouse and rat origin and MAGOHB of mouse origin by Western Blotting (starting dilution 1:200, dilution range 1:100-1:1000), immunofluorescence (starting dilution 1:50, dilution range 1:50-1:500) and solid phase ELISA (starting dilution 1:30, dilution range 1:30-1:3000).

MAGOH (C-12) is also recommended for detection of MAGOH in additional species, including equine, canine, bovine, porcine and avian.

Suitable for use as control antibody for MAGOH siRNA (h): sc-60978, MAGOH shRNA Plasmid (h): sc-60978-SH and MAGOH shRNA (h) Lentiviral Particles: sc-60978-V.

MAGOH (C-12) X TransCruz antibody is recommended for Gel Supershift and ChIP applications.

Molecular Weight of MAGOH: 17 kDa.

Positive Controls: Ramos cell lysate: sc-2216, Ramos nuclear extract: sc-2153 or BJAB nuclear extract: sc-2145.

RECOMMENDED SECONDARY REAGENTS

To ensure optimal results, the following support (secondary) reagents are recommended: 1) Western Blotting: use donkey anti-goat IgG-HRP: sc-2020 (dilution range: 1:2000-1:100,000) or Cruz Marker™ compatible donkey anti-goat IgG-HRP: sc-2033 (dilution range: 1:2000-1:5000), Cruz Marker™ Molecular Weight Standards: sc-2035, TBS Blotto A Blocking Reagent: sc-2333 and Western Blotting Luminol Reagent: sc-2048. 2) Immunofluorescence: use donkey anti-goat IgG-FITC: sc-2024 (dilution range: 1:100-1:400) or donkey anti-goat IgG-TR: sc-2783 (dilution range: 1:100-1:400) with UltraCruz™ Mounting Medium: sc-24941.

STORAGE

Store at 4° C, **DO NOT FREEZE**. Stable for one year from the date of shipment. Non-hazardous. No MSDS required.

PROTOCOLS

See our web site at www.scbt.com or our catalog for detailed protocols and support products.



Try **MAGOH (21B12): sc-56724** or **MAGOH (F-6): sc-271365**, our highly recommended monoclonal alternatives to MAGOH (C-12).