

PC-PLD1 (G-13): sc-48258

BACKGROUND

Virtually every cell uses phosphatidylcholine as a substrate to produce phosphatidic acid and choline. Phosphatidylcholine phospholipase D1 and D2 (PC-PLD1 and PC-PLD2) are phospholipid-specific phosphodiesterases that hydrolyze phosphatidylcholine. Unlike PC-PLD1, which associates with secretory granules, PC-PLD2 localizes to the plasma membrane, where it is implicated in the formation of endocytotic vesicles. Both PC-PLD1 and PC-PLD2 coordinately regulate macrophage phagocytosis. PC-PLD activity in mammalian cells is transiently stimulated upon activation by G protein-coupled and receptor tyrosine kinase cell surface receptors. For example, PC-PLD1 and PC-PLD2 participate in sphingosine 1-phosphate stimulation of ERK phosphorylation and IL-8 secretion in bronchial epithelial cells. In addition, tubulin binding to PC-PLD2 inhibits muscarinic receptor-linked PC-PLD2 activation. PC-PLD2 also enhances PKC ζ activity through direct interaction in a lipase activity-independent manner. PC-PLD1 and PC-PLD2 stimulate cell growth by repressing expression of p21 gene through p53-dependent and p53-independent pathways, respectively, which may ultimately lead to carcinogenesis.

REFERENCES

1. Nishida, A., et al. 1994. Brain ischemia decreases phosphatidylcholine-phospholipase D but not phosphatidylinositol-phospholipase C in rats. *Stroke* 25: 1247-1251.
2. del Peso, L., et al. 1996. Activation of phospholipase D by Ras proteins is independent of protein kinase C. *J. Cell. Biochem.* 61: 599-608.
3. Houle, M.G., et al. 1999. Regulation of phospholipase D by phosphorylation-dependent mechanisms. *Biochim. Biophys. Acta* 1439: 135-149.
4. Cockcroft, S. 2001. Signalling roles of mammalian phospholipase D1 and D2. *Cell. Mol. Life Sci.* 58: 1674-1687.
5. Zhao, D., et al. 2001. Generation of choline for acetylcholine synthesis by phospholipase D isoforms. *BMC Neurosci.* 2: 16.
6. Chahdi, A., et al. 2002. Serine/threonine protein kinases synergistically regulate phospholipase D1 and 2 and secretion in RBL-2H3 mast cells. *Mol. Immunol.* 38:1269-1276.

CHROMOSOMAL LOCATION

Genetic locus: PLD1 (human) mapping to 3q26.31; Pld1 (mouse) mapping to 3 A3.

SOURCE

PC-PLD1 (G-13) is an affinity purified goat polyclonal antibody raised against a peptide mapping within an internal region of PC-PLD1 of human origin.

PRODUCT

Each vial contains 200 μ g IgG in 1.0 ml of PBS with < 0.1% sodium azide and 0.1% gelatin.

Blocking peptide available for competition studies, sc-48258 P, (100 μ g peptide in 0.5 ml PBS containing < 0.1% sodium azide and 0.2% BSA).

APPLICATIONS

PC-PLD1 (G-13) is recommended for detection of PC-PLD1 of mouse, rat and human origin by Western Blotting (starting dilution 1:200, dilution range 1:100-1:1000), immunofluorescence (starting dilution 1:50, dilution range 1:50-1:500) and solid phase ELISA (starting dilution 1:30, dilution range 1:30-1:3000).

PC-PLD1 (G-13) is also recommended for detection of PC-PLD1 in additional species, including equine, canine, bovine and porcine.

Suitable for use as control antibody for PC-PLD1 siRNA (h): sc-44000, PC-PLD1 siRNA (m): sc-41629, PC-PLD1 shRNA Plasmid (h): sc-44000-SH, PC-PLD1 shRNA Plasmid (m): sc-41629-SH, PC-PLD1 shRNA (h) Lentiviral Particles: sc-44000-V and PC-PLD1 shRNA (m) Lentiviral Particles: sc-41629-V.

Molecular Weight of PC-PLD1 α : 120 kDa.

Molecular Weight of PC-PLD1 β : 115 kDa.

Positive Controls: MCF7 + insulin cell lysate: sc-24733, Caki-1 cell lysate: sc-2224 or HeLa whole cell lysate: sc-2200.

RECOMMENDED SECONDARY REAGENTS

To ensure optimal results, the following support (secondary) reagents are recommended: 1) Western Blotting: use donkey anti-goat IgG-HRP: sc-2020 (dilution range: 1:2000-1:100,000) or Cruz Marker[™] compatible donkey anti-goat IgG-HRP: sc-2033 (dilution range: 1:2000-1:5000), Cruz Marker[™] Molecular Weight Standards: sc-2035, TBS Blotto A Blocking Reagent: sc-2333 and Western Blotting Luminol Reagent: sc-2048. 2) Immunofluorescence: use donkey anti-goat IgG-FITC: sc-2024 (dilution range: 1:100-1:400) or donkey anti-goat IgG-TR: sc-2783 (dilution range: 1:100-1:400) with UltraCruz[™] Mounting Medium: sc-24941.

STORAGE

Store at 4° C, ****DO NOT FREEZE****. Stable for one year from the date of shipment. Non-hazardous. No MSDS required.

RESEARCH USE

For research use only, not for use in diagnostic procedures.

PROTOCOLS

See our web site at www.scbt.com or our catalog for detailed protocols and support products.


 MONOS
 Satisfaction
 Guaranteed

Try **PC-PLD1 (F-12): sc-28314**, our highly recommended monoclonal alternative to PC-PLD1 (G-13).