

# RGS20 (P-12): sc-48295

## BACKGROUND

The regulators of G protein signaling (RGS) proteins inhibit heterotrimeric G protein signaling. RGS proteins work by functioning as GTPase-activating proteins (which increase the GTPase activity of G protein  $\alpha$  subunits) thereby driving G proteins into their inactive GDP-bound form. RGS20 is a 25 kDa protein expressed exclusively in brain, with highest levels in the caudate nucleus and temporal lobe. RGS20 belongs to the RZ subfamily because it is highly selective for the  $\alpha_z$  subunit on G proteins. However, if protein kinase C phosphorylates the  $\alpha_z$  subunit, the G protein is much less susceptible to RGS20 action. RGS20 directly interacts with the microtubule-destabilizing protein SCG10 (superior cervical ganglia, neural specific 10) and blocks its ability to induce microtubule disassembly.

## REFERENCES

- Glick, J.L., et al. 1998. RGSZ1, a  $G_z$ -selective regulator of G protein signaling whose action is sensitive to the phosphorylation state of  $G_{\alpha_z}$ . *J. Biol. Chem.* 273: 26008-26013.
- Wang, J., et al. 1998. RGSZ1, a  $G_z$ -selective RGS protein in brain. Structure, membrane association, regulation by  $G_{\alpha_z}$  phosphorylation and relationship to a  $G_z$  GTPase-activating protein subfamily. *J. Biol. Chem.* 273: 26014-26025.
- Barker, S.A., et al. 2001. RGSZ1 and Ret RGS: two of several splice variants from the gene RGS20. *Genomics* 78: 223-229.
- Nixon, A.B., et al. 2002. The interaction of RGSZ1 with SCG10 attenuates the ability of SCG10 to promote microtubule disassembly. *J. Biol. Chem.* 277: 18127-18133.
- Online Mendelian Inheritance in Man, OMIM™. 2002. Johns Hopkins University, Baltimore, MD. MIM Number: 607190. World Wide Web URL: <http://www.ncbi.nlm.nih.gov/omim/>
- Wang, Y., et al. 2004. Analysis of RGSZ1 protein interaction with  $G_{\alpha_i}$  subunits. *Meth. Enzymol.* 390: 31-52.
- Nixon, A.B., et al. 2004. Analysis of the regulation of microtubule dynamics by interaction of RGSZ1 (RGS20) with the neuronal stathmin, SCG10. *Meth. Enzymol.* 390: 53-64.

## CHROMOSOMAL LOCATION

Genetic locus: RGS20 (human) mapping to 8q11.23; Rgs20 (mouse) mapping to 1 A1.

## SOURCE

RGS20 (P-12) is an affinity purified goat polyclonal antibody raised against a peptide mapping at the C-terminus of RGS20 of human origin.

## PRODUCT

Each vial contains 200  $\mu$ g IgG in 1.0 ml of PBS with < 0.1% sodium azide and 0.1% gelatin.

Blocking peptide available for competition studies, sc-48295 P, (100  $\mu$ g peptide in 0.5 ml PBS containing < 0.1% sodium azide and 0.2% BSA).

## APPLICATIONS

RGS20 (P-12) is recommended for detection of all isoforms of RGS20 of mouse, rat and human origin by Western Blotting (starting dilution 1:200, dilution range 1:100-1:1000), immunoprecipitation [1-2  $\mu$ g per 100-500  $\mu$ g of total protein (1 ml of cell lysate)], immunofluorescence (starting dilution 1:50, dilution range 1:50-1:500) and solid phase ELISA (starting dilution 1:30, dilution range 1:30-1:3000).

RGS20 (P-12) is also recommended for detection of all isoforms of RGS20 in additional species, including equine, canine, bovine and avian.

Suitable for use as control antibody for RGS20 siRNA (h): sc-61470, RGS20 siRNA (m): sc-61471, RGS20 shRNA Plasmid (h): sc-61470-SH, RGS20 shRNA Plasmid (m): sc-61471-SH, RGS20 shRNA (h) Lentiviral Particles: sc-61470-V and RGS20 shRNA (m) Lentiviral Particles: sc-61471-V.


Molecular Weight of RGS20: 44 kDa.

Positive Controls: RGS20 (h): 293T Lysate: sc-372921, SK-N-SH cell lysate: sc-2410 or HeLa whole cell lysate: sc-2200.

## RECOMMENDED SECONDARY REAGENTS

To ensure optimal results, the following support (secondary) reagents are recommended: 1) Western Blotting: use donkey anti-goat IgG-HRP: sc-2020 (dilution range: 1:2000-1:100,000) or Cruz Marker™ compatible donkey anti-goat IgG-HRP: sc-2033 (dilution range: 1:2000-1:5000), Cruz Marker™ Molecular Weight Standards: sc-2035, TBS Blotto A Blocking Reagent: sc-2333 and Western Blotting Luminol Reagent: sc-2048. 2) Immunoprecipitation: use Protein A/G PLUS-Agarose: sc-2003 (0.5 ml agarose/2.0 ml). 3) Immunofluorescence: use donkey anti-goat IgG-FITC: sc-2024 (dilution range: 1:100-1:400) or donkey anti-goat IgG-TR: sc-2783 (dilution range: 1:100-1:400) with UltraCruz™ Mounting Medium: sc-24941.

## DATA



RGS20 (P-12): sc-48295. Western blot analysis of RGS20 expression in non-transfected: sc-117752 (A) and human RGS20 transfected: sc-372921 (B) 293T whole cell lysates.

## STORAGE

Store at 4° C, \*\*DO NOT FREEZE\*\*. Stable for one year from the date of shipment. Non-hazardous. No MSDS required.

## RESEARCH USE

For research use only, not for use in diagnostic procedures.