

p40-phox (B-1): sc-48376



The Power to Question

BACKGROUND

Nicotinamide adenine dinucleotide phosphate (NADPH)-oxidase is a multi-meric enzyme system that mediates electron transport from NADPH in the cytoplasm to molecular oxygen in the phagosome, thereby generating reactive oxidant intermediates. Upon neutrophil stimulation, NADPH-oxidase and other cytosolic elements localize to the cell membrane from the cytosol to form a complex which produces phagocytic oxygen radicals. There are a number of cytosolic proteins that are involved in NADPH-oxidase activation/deactivation, including p47-phox, p67-phox, p40-phox and the small GTP-binding protein, Rac. Activation of NADPH oxidase is accompanied by the phosphorylation of cytosolic components p40-phox, p47-phox and p67-phox. The PKC consensus phosphorylation sites Thr 154 and Ser 315 in p40-phox are phosphorylated during activation of NADPH oxidase. p40-phox can promote oxidase activation by increasing the affinity of p47-phox for NADPH-oxidase. However, p40-phox appears to downregulate oxidase function as well, by competing with an SH3 domain interaction between other essential oxidase components.

CHROMOSOMAL LOCATION

Genetic locus: NCF4 (human) mapping to 22q12.3; Ncf4 (mouse) mapping to 15 E1.

SOURCE

p40-phox (B-1) is a mouse monoclonal antibody raised against amino acids 1-300 p40-phox of human origin.

PRODUCT

Each vial contains 200 µg IgG_{2b} kappa light chain in 1.0 ml of PBS with < 0.1% sodium azide and 0.1% gelatin.

STORAGE

Store at 4° C, **DO NOT FREEZE**. Stable for one year from the date of shipment. Non-hazardous. No MSDS required.

RESEARCH USE

For research use only, not for use in diagnostic procedures.

APPLICATIONS

p40-phox (B-1) is recommended for detection of p40-phox of mouse, rat and human origin by Western Blotting (starting dilution 1:100, dilution range 1:100-1:1000), immunoprecipitation [1-2 µg per 100-500 µg of total protein (1 ml of cell lysate)], immunofluorescence (starting dilution 1:50, dilution range 1:50-1:500) and solid phase ELISA (starting dilution 1:30, dilution range 1:30-1:3000).

Suitable for use as control antibody for p40-phox siRNA (h): sc-36155, p40-phox siRNA (m): sc-36156, p40-phox shRNA Plasmid (h): sc-36155-SH, p40-phox shRNA Plasmid (m): sc-36156-SH, p40-phox shRNA (h) Lentiviral Particles: sc-36155-V and p40-phox shRNA (m) Lentiviral Particles: sc-36156-V.


Molecular Weight of p40-phox: 40 kDa.

Positive Controls: THP-1 cell lysate: sc-2238, HL-60 whole cell lysate: sc-2209 or HEL 92.1.7 cell lysate: sc-2270.


RECOMMENDED SUPPORT REAGENTS

To ensure optimal results, the following support reagents are recommended: 1) Western Blotting: use m-IgGκ BP-HRP: sc-516102 or m-IgGκ BP-HRP (Cruz Marker): sc-516102-CM (dilution range: 1:1000-1:10000), Cruz Marker™ Molecular Weight Standards: sc-2035, UltraCruz® Blocking Reagent: sc-516214 and Western Blotting Luminol Reagent: sc-2048. 2) Immunoprecipitation: use Protein A/G PLUS-Agarose: sc-2003 (0.5 ml agarose/2.0 ml). 3) Immunofluorescence: use m-IgGκ BP-FITC: sc-516140 or m-IgGκ BP-PE: sc-516141 (dilution range: 1:50-1:200) with UltraCruz® Mounting Medium: sc-24941 or UltraCruz® Hard-set Mounting Medium: sc-359850.

DATA



p40-phox (B-1): sc-48376. Western blot analysis of p40-phox expression in THP-1 (A), HL-60 (B), HEL 92.1.7 (C), RPMI-8226 (D), U-937 (E) and U-698-M (F) whole cell lysates. Detection reagent used: m-IgG_{2b} BP-HRP: sc-542741.



Western blot analysis of p40-phox phosphorylation in untreated (A, C) and lambda protein phosphatase (sc-200312A) treated (B, D) THP-1 whole cell lysates. Antibodies tested include p-p40-phox (Thr 154): sc-33403 (A, B) and p40-phox (B-1): sc-48376 (C, D).

SELECT PRODUCT CITATIONS

- Chen, L., et al. 2011. Cadmium induction of reactive oxygen species activates the mTOR pathway, leading to neuronal cell death. *Free Radic. Biol. Med.* 50: 624-632.
- Ansari, M.A., et al. 2014. A time course of NADPH-oxidase up-regulation and endothelial nitric oxide synthase activation in the hippocampus following neurotrauma. *Free Radic. Biol. Med.* 77: 21-29.
- Gluschko, A., et al. 2022. Macrophages target *Listeria monocytogenes* by two discrete non-canonical autophagy pathways. *Autophagy* 18: 1090-1107.
- Tlili, A., et al. 2023. ROCK2 interacts with p22phox to phosphorylate p47phox and to control NADPH oxidase activation in human monocytes. *Proc. Natl. Acad. Sci. USA* 120: e2209184120.

PROTOCOLS

See our web site at www.scbt.com for detailed protocols and support products.