

GPR30 (N-15)-R: sc-48525-R

BACKGROUND

G protein-coupled receptors (GPCRs), also designated seven transmembrane (7TM) receptors and heptahelical receptors, are a protein family which interact with G proteins (heterotrimeric GTPases) to synthesize intracellular second messengers such as diacylglycerol, cyclic AMP, inositol phosphates and calcium ions. Their diverse biological functions range from vision and olfaction to neuronal and endocrine signaling and are involved in many pathological conditions. G protein receptor 30 (GPR30), also designated chemokine receptor-like 2 (CMKRL2), is a 375 amino acid protein orphan GPCR. GPR30 is an intracellular transmembrane estrogen receptor localized to the endoplasmic reticulum which binds estrogen and estrogen derivatives.

CHROMOSOMAL LOCATION

Genetic locus: GPER (human) mapping to 7p22.3; Gper (mouse) mapping to 5 G2.

SOURCE

GPR30 (N-15)-R is an affinity purified rabbit polyclonal antibody raised against a peptide mapping near the N-terminus of GPR30 of human origin.

PRODUCT

Each vial contains 200 µg IgG in 1.0 ml of PBS with < 0.1% sodium azide and 0.1% gelatin.

Blocking peptide available for competition studies, sc-48525 P, (100 µg peptide in 0.5 ml PBS containing < 0.1% sodium azide and 0.2% BSA).

APPLICATIONS

GPR30 (N-15)-R is recommended for detection of GPR30 of mouse, rat and human origin by Western Blotting (starting dilution 1:200, dilution range 1:100-1:1000), immunoprecipitation [1-2 µg per 100-500 µg of total protein (1 ml of cell lysate)], immunofluorescence (starting dilution 1:50, dilution range 1:50-1:500), immunohistochemistry (including paraffin-embedded sections) (starting dilution 1:50, dilution range 1:50-1:500) and solid phase ELISA (starting dilution 1:30, dilution range 1:30-1:3000).

Suitable for use as control antibody for GPR30 siRNA (h): sc-60743, GPR30 siRNA (m): sc-60744, GPR30 siRNA (r): sc-156143, GPR30 shRNA Plasmid (h): sc-60743-SH, GPR30 shRNA Plasmid (m): sc-60744-SH, GPR30 shRNA Plasmid (r): sc-156143-SH, GPR30 shRNA (h) Lentiviral Particles: sc-60743-V, GPR30 shRNA (m) Lentiviral Particles: sc-60744-V, and GPR30 shRNA (r) Lentiviral Particles: sc-156143-V.

Molecular Weight of GPR30: 38 kDa.

Positive Controls: HeLa whole cell lysate: sc-2200, MCF7 whole cell lysate: sc-2206 or JAR cell lysate: sc-2276.


STORAGE

Store at 4° C, ****DO NOT FREEZE****. Stable for one year from the date of shipment. Non-hazardous. No MSDS required.


RESEARCH USE

For research use only, not for use in diagnostic procedures.

DATA



GPR30 (N-15)-R: sc-48525-R. Western blot analysis of GPR30 expression in SK-BR-3 (A), HeLa (B), ZR-75-1 (C), MCF7 (D) and JAR (E) whole cell lysates.



GPR30 (N-15)-R: sc-48525-R. Immunofluorescence staining of methanol-fixed HeLa cells showing membrane localization.

SELECT PRODUCT CITATIONS

- Miñano, A., et al. 2008. Estradiol facilitates neurite maintenance by a Src/Ras/ERK signalling pathway. *Mol. Cell. Neurosci.* 39: 143-151.
- Pupo, M., et al. 2013. The nuclear localization signal is required for nuclear GPER translocation and function in breast cancer-associated fibroblasts (CAFs). *Mol. Cell. Endocrinol.* 376: 23-32.
- De Francesco, E.M., et al. 2013. GPER mediates cardiotropic effects in spontaneously hypertensive rat hearts. *PLoS ONE* 8: e69322.
- Santolla, M.F., et al. 2014. Niacin activates the G protein-coupled estrogen receptor (GPER)-mediated signalling. *Cell. Signal.* 26: 1466-1475.
- De Francesco, E.M., et al. 2014. GPER mediates activation of HIF1 α /VEGF signaling by estrogens. *Cancer Res.* 74: 4053-4064.
- De Marco, P., et al. 2014. GPER1 is regulated by insulin in cancer cells and cancer-associated fibroblasts. *Endocr. Relat. Cancer* 21: 739-753.
- Pellegrini, M., et al. 2014. Endocrine disruptors differently influence estrogen receptor β and androgen receptor in male and female rat VSMC. *J. Cell. Physiol.* 229: 1061-1068.
- Comeglio, P., et al. 2014. Opposite effects of tamoxifen on metabolic syndrome-induced bladder and prostate alterations: a role for GPR30/GPER? *Prostate* 74: 10-28.
- Albanito, L., et al. 2015. Effects of atrazine on estrogen receptor α - and G protein-coupled receptor 30-mediated signaling and proliferation in cancer cells and cancer-associated fibroblasts. *Environ. Health Perspect.* 123: 493-499.
- Sinicropi, M.S., et al. 2015. (6-Bromo-1,4-dimethyl-9H-carbazol-3-yl-methylene)-hydrazine (Carbhydraz) acts as a GPER agonist in breast cancer cells. *Curr. Top. Med. Chem.* 15: 1035-1042.
- Vivacqua, A., et al. 2015. Estrogenic gper signaling regulates mir144 expression in cancer cells and cancer-associated fibroblasts. *Oncotarget* 6: 16573-16587.
- Rigiracciolo, D.C., et al. 2015. Copper activates HIF-1 α /GPER/VEGF signalling in cancer cells. *Oncotarget* 6: 34158-34177.