

KIF12 (N-18): sc-48558

BACKGROUND

Kinesin is a cytoskeletal motor protein involved in axonal transport and cell division. The kinesin superfamily proteins (KIFs) are microtubule-dependent molecular motors that transport membranous organelles and protein complexes in a microtubule- and ATP-dependent manner. Cells use KIFs to tightly control the direction, destination, and speed of transportation of a variety of important functional molecules, including mRNA. KIFs are involved in neuronal function and development. Kinesin family member 12 (KIF12) is a kinesin-like 651-amino-acid protein which is involved in mitotically linked cytokinesis. KIF12 is required during mitosis for normal myosin II localization and during late anaphase and telophase for normal nuclear separation. The KIF12 gene consists of a KISc domain, a coiled-coil domain with an internal hinge region and a C-terminal tail domain. KIF12 mRNA is expressed in fetal liver, adult brain, pancreatic islet, kidney tumors, and uterine and pancreatic cancers.

REFERENCES

- Howard, J. 1996. The movement of kinesin along microtubules. *Annu. Rev. Physiol.* 58: 703-729.
- Miki, H., et al. 2001. All kinesin superfamily protein, KIF, genes in mouse and human. *Proc. Natl. Acad. Sci. USA* 98: 7004-7011.
- Mburu, P., et al. 2003. Defects in Whirlin, a PDZ domain molecule involved in stereocilia elongation, cause deafness in the whirler mouse and families with DFNB31. *Nat. Genet.* 34: 421-428.
- Lakshminanth, G.S., et al. 2004. A mitotic kinesin-like protein required for normal karyokinesis, myosin localization to the furrow, and cytokinesis in *Dictyostelium*. *Proc. Natl. Acad. Sci. USA* 101: 16519-16524.
- Mrug, M., et al. 2005. Kinesin family member 12 is a candidate polycystic kidney disease modifier in the cpk mouse. *J. Am. Soc. Nephrol.* 16: 905-916.

CHROMOSOMAL LOCATION

Genetic locus: KIF12 (human) mapping to 9q32; Kif12 (mouse) mapping to 4 B3.

SOURCE

KIF12 (N-18) is an affinity purified goat polyclonal antibody raised against a peptide mapping near the N-terminus of KIF12 of human origin.

PRODUCT

Each vial contains 200 µg IgG in 1.0 ml of PBS with < 0.1% sodium azide and 0.1% gelatin.

Blocking peptide available for competition studies, sc-48558 P, (100 µg peptide in 0.5 ml PBS containing < 0.1% sodium azide and 0.2% BSA).

STORAGE

Store at 4° C, ****DO NOT FREEZE****. Stable for one year from the date of shipment. Non-hazardous. No MSDS required.

RESEARCH USE

For research use only, not for use in diagnostic procedures.

APPLICATION

KIF12 (N-18) is recommended for detection of KIF12 of mouse and human origin by Western Blotting (starting dilution 1:200, dilution range 1:100-1:1000), immunoprecipitation [1-2 µg per 100-500 µg of total protein (1 ml of cell lysate)], immunofluorescence (starting dilution 1:50, dilution range 1:50-1:500) and solid phase ELISA (starting dilution 1:30, dilution range 1:30-1:3000).

KIF12 (N-18) is also recommended for detection of KIF12 in additional species, including equine, canine and porcine.


Suitable for use as control antibody for KIF12 siRNA (h): sc-60880, KIF12 siRNA (m): sc-60881, KIF12 shRNA Plasmid (h): sc-60880-SH, KIF12 shRNA Plasmid (m): sc-60881-SH, KIF12 shRNA (h) Lentiviral Particles: sc-60880-V and KIF12 shRNA (m) Lentiviral Particles: sc-60881-V.

Molecular Weight (predicted) of KIF12: 70 kDa.

Molecular Weight (observed) of KIF12: 70-85 kDa.

Positive Controls: KIF12 (h): 293T Lysate: sc-111091 or MIA PaCa-2 cell lysate: sc-2285.

DATA



KIF12 (N-18): sc-48558. Western blot analysis of KIF12 expression in non-transfected: sc-117752 (A) and truncated human KIF12 transfected: sc-111091 (B) 293T whole cell lysates.

PROTOCOLS

See our web site at www.scbt.com or our catalog for detailed protocols and support products.


 MONOS
Satisfaction
Guaranteed

Try **KIF12 (C-4): sc-376572** or **KIF12 (A-9): sc-376766**, our highly recommended monoclonal alternatives to KIF12 (N-18).