

# MATP (N-14): sc-48592

## BACKGROUND

The membrane-associated transporter protein (MATP) is a 530 amino acid protein that spans the lipid bilayer 12 times. MATP is a melanocyte differentiation antigen that is expressed in a high percentage of melanoma cell lines. MATP is transcriptionally modulated by MITF, a melanocyte-specific transcription factor that may act indirectly or bind to a remote regulatory sequence. MATP may play a role in skin cancer, as its gene is expressed in a high percentage of melanoma cell lines, but not at significant levels in normal tissues. Mutations in the MATP gene have also been linked to albinism.

## REFERENCES

1. Baxter, L.L. and Pavan, W.J. 2002. The oculocutaneous albinism type IV gene MATP is a new marker of pigment cell precursors during mouse embryonic development. *Mech. Dev.* 116: 209-212.
2. Online Mendelian Inheritance in Man, OMIM™. 2002. Johns Hopkins University, Baltimore, MD. MIM Number: 606202. World Wide Web URL: <http://www.ncbi.nlm.nih.gov/omim/>
3. Rundshagen, U., Zühlke, C., Opitz, S., Schwinger, E. and Käsmann-Kellner, B. 2004. Mutations in the MATP gene in five German patients affected by oculocutaneous albinism type 4. *Hum. Mutat.* 23: 106-110.
4. Yuasa, I., Umetsu, K., Watanabe, G., Nakamura, H., Endoh, M. and Irizawa, Y. 2004. MATP polymorphisms in Germans and Japanese: the L374F mutation as a population marker for Caucasoids. *Int. J. Legal Med.* 118: 364-366.
5. Blalock, J.E. 2005. The immune system as the sixth sense. *J. Intern. Med.* 257: 126-138.
6. Graf, J., Hodgson, R. and van Daal, A. 2005. Single nucleotide polymorphisms in the MATP gene are associated with normal human pigmentation variation. *Hum. Mutat.* 25: 278-284.
7. Inagaki, K., Suzuki, T., Ito, S., Suzuki, N., Fukai, K., Horiuchi, T., Tanaka, T., Manabe, E. and Tomita, Y. 2005. OCA4: evidence for a founder effect for the p.D157N mutation of the MATP gene in Japanese and Korean. *Pigment Cell Res.* 18: 385-388.
8. Suzuki, T., Inagaki, K., Fukai, K., Obana, A., Lee, S.T. and Tomita, Y. 2005. A Korean case of oculocutaneous albinism type IV caused by a D157N mutation in the MATP gene. *Br. J. Dermatol.* 152: 174-175.

## CHROMOSOMAL LOCATION

Genetic locus: MATP (human) mapping to 5p13.3; Matp (mouse) mapping to 15 A1.

## SOURCE

MATP (N-14) is an affinity purified goat polyclonal antibody raised against a peptide mapping at the N-terminus of MATP of human origin.

## STORAGE

Store at 4° C, **\*\*DO NOT FREEZE\*\***. Stable for one year from the date of shipment. Non-hazardous. No MSDS required.

## PRODUCT

Each vial contains 200 µg IgG in 1.0 ml of PBS with < 0.1% sodium azide and 0.1% gelatin.

Blocking peptide available for competition studies, sc-48592 P, (100 µg peptide in 0.5 ml PBS containing < 0.1% sodium azide and 0.2% BSA).

## APPLICATIONS

MATP (N-14) is recommended for detection of MATP of mouse, rat and human origin by Western Blotting (starting dilution 1:200, dilution range 1:100-1:1000), immunofluorescence (starting dilution 1:50, dilution range 1:50-1:500) and solid phase ELISA (starting dilution 1:30, dilution range 1:30-1:3000).

Suitable for use as control antibody for MATP siRNA (h): sc-60986, MATP siRNA (m): sc-60987, MATP shRNA Plasmid (h): sc-60986-SH, MATP shRNA Plasmid (m): sc-60987-SH, MATP shRNA (h) Lentiviral Particles: sc-60986-V and MATP shRNA (m) Lentiviral Particles: sc-60987-V.

Molecular Weight of MATP: 58 kDa.

## RECOMMENDED SECONDARY REAGENTS

To ensure optimal results, the following support (secondary) reagents are recommended: 1) Western Blotting: use donkey anti-goat IgG-HRP: sc-2020 (dilution range: 1:2000-1:100,000) or Cruz Marker™ compatible donkey anti-goat IgG-HRP: sc-2033 (dilution range: 1:2000-1:5000), Cruz Marker™ Molecular Weight Standards: sc-2035, TBS Blotto A Blocking Reagent: sc-2333 and Western Blotting Luminol Reagent: sc-2048. 2) Immunofluorescence: use donkey anti-goat IgG-FITC: sc-2024 (dilution range: 1:100-1:400) or donkey anti-goat IgG-TR: sc-2783 (dilution range: 1:100-1:400) with UltraCruz™ Mounting Medium: sc-24941.

## RESEARCH USE

For research use only, not for use in diagnostic procedures.

## PROTOCOLS

See our web site at [www.scbt.com](http://www.scbt.com) or our catalog for detailed protocols and support products.


 MONOS  
 Satisfaction  
 Guaranteed

Try **MATP (F-4): sc-377397** or **MATP (7K-2): sc-100780**, our highly recommended monoclonal alternatives to MATP (N-14).