

DSC3 (H-50): sc-48750

BACKGROUND

The desmosomal cadherin desmocollin DSC1 is expressed in upper epidermis where strong adhesion is required. DSC1 is a type I membrane protein required for strong adhesion and barrier maintenance in epidermis and contributes to epidermal differentiation. DSC3 is also a type I membrane protein and is expressed in all living epidermal layers as well as in glandular ducts and in basal matrix cells and the outer root sheath of hair follicles. DSC3, but not DSC1, is also present in desmosomes of the basal and suprabasal cell layers of other stratified epithelia such as cervix, tongue and esophagus as well as in cells of the basal layer of bladder urothelium and the complex epithelium of trachea. The DSC1 gene comprises 17 exons spanning approximately 33 kb on 18q12.1, and the DSC3 gene comprises 17 exons spanning approximately 49 kb on 18q12.1. Alternative splicing gives rise to two DSC1 and DSC3 isoforms, designated 1A and 1B, and 3A and 3B, respectively, which differ in their carboxy termini. DSC1 and DSC3 may contribute to epidermal cell positioning by mediating differential adhesiveness between cells that express different isoforms.

CHROMOSOMAL LOCATION

Genetic locus: DSC3 (human) mapping to 18q12.1; DSC3 (mouse) mapping to 18 A2.

SOURCE

DSC3 (H-50) is a rabbit polyclonal antibody raised against amino acids 641-690 mapping within an extracellular domain of DSC3 of human origin.

PRODUCT

Each vial contains 200 µg IgG in 1.0 ml of PBS with < 0.1% sodium azide and 0.1% gelatin.

STORAGE

Store at 4° C, ****DO NOT FREEZE****. Stable for one year from the date of shipment. Non-hazardous. No MSDS required.

APPLICATIONS

DSC3 (H-50) is recommended for detection of DSC3A and DSC3B of mouse and human origin by Western Blotting (starting dilution 1:200, dilution range 1:100-1:1000), immunoprecipitation [1-2 µg per 100-500 µg of total protein (1 ml of cell lysate)], immunofluorescence (starting dilution 1:50, dilution range 1:50-1:500) and solid phase ELISA (starting dilution 1:30, dilution range 1:30-1:3000).

Suitable for use as control antibody for DSC3 siRNA (h): sc-43111, DSC3 siRNA (m): sc-43112, DSC3 shRNA Plasmid (h): sc-43111-SH, DSC3 shRNA Plasmid (m): sc-43112-SH, DSC3 shRNA (h) Lentiviral Particles: sc-43111-V and DSC3 shRNA (m) Lentiviral Particles: sc-43112-V.


Molecular Weight of DSC3: 100 kDa.

Positive Controls: DSC3 (m): 293 Lysate: sc-178531, FHs 173We cell lysate: sc-2417 or Jurkat whole cell lysate: sc-2204.


RECOMMENDED SECONDARY REAGENTS

To ensure optimal results, the following support (secondary) reagents are recommended: 1) Western Blotting: use goat anti-rabbit IgG-HRP: sc-2004 (dilution range: 1:2000-1:100,000) or Cruz Marker™ compatible goat anti-rabbit IgG-HRP: sc-2030 (dilution range: 1:2000-1:5000), Cruz Marker™ Molecular Weight Standards: sc-2035, TBS Blotto A Blocking Reagent: sc-2333 and Western Blotting Luminol Reagent: sc-2048. 2) Immunoprecipitation: use Protein A/G PLUS-Agarose: sc-2003 (0.5 ml agarose/2.0 ml). 3) Immunofluorescence: use goat anti-rabbit IgG-FITC: sc-2012 (dilution range: 1:100-1:400) or goat anti-rabbit IgG-TR: sc-2780 (dilution range: 1:100-1:400) with UltraCruz™ Mounting Medium: sc-24941.

DATA



DSC3 (H-50): sc-48750. Western blot analysis of DSC3 expression in non-transfected 293: sc-110760 (A), mouse DSC3 transfected 293: sc-178531 (B), FHs 173We (C) and NIH/3T3 (D) whole cell lysates.



DSC3 (H-50): sc-48750. Western blot analysis of DSC3 expression in THP-1 (A) and Jurkat (B) whole cell lysates.

SELECT PRODUCT CITATIONS

- Mao, X., Choi, E.J. and Payne, A.S. 2009. Disruption of desmosome assembly by monovalent human pemphigus vulgaris monoclonal antibodies. *J. Invest. Dermatol.* 129: 908-918.

RESEARCH USE

For research use only, not for use in diagnostic procedures.

PROTOCOLS

See our web site at www.scbt.com or our catalog for detailed protocols and support products.

MONOS
Satisfaction
Guaranteed

Try **DSC2/3 (7G6): sc-53485** or **DSC3 (93-71): sc-81806**, our highly recommended monoclonal alternatives to DSC3 (H-50).