

# TREM-2 (H-160): sc-48764

## BACKGROUND

Polycystic lipomembranous osteodysplasia with sclerosing leukoencephalopathy (PLOS), known as "Nasu-Hakola disease", is a recessively inherited disease where individuals display early-onset progressive dementia and bone cysts, which leads to death. Mutations in TYROBP (DAP12), which codes for a membrane receptor component in natural-killer and myeloid cells and mutations in triggering receptor expressed on myeloid cells-2 (TREM-2), correlate well to the pathology of PLOS. TREM-2 is a cell surface receptor on human monocyte-derived dendritic cells that forms a receptor signaling complex with DAP12 and triggers activation of the immune response in macrophages and dendritic cells (DC). The TREM-2/DAP12 complex is a molecular promoter of upregulation of CC chemokine receptor 7, partial DC maturation and DC survival through activation of protein tyrosine kinases and extracellular signal-regulated kinase. The human chronic inflammatory TREM-2 gene maps to chromosome 6p21.1 and encodes a 230 amino acid protein.

## REFERENCES

1. Bouchon, A., et al. 2001. A DAP12-mediated pathway regulates expression of CC chemokine receptor 7 and maturation of human dendritic cells. *J. Exp. Med.* 194: 1111-1122.
2. Daws, M.R., et al. 2001. Cloning and characterization of a novel mouse myeloid DAP12-associated receptor family. *Eur. J. Immunol.* 31: 783-791.
3. Paloneva, J., et al. 2002. Mutations in two genes encoding different subunits of a receptor signaling complex result in an identical disease phenotype. *Am. J. Hum. Genet.* 71: 656-662.
4. Online Mendelian Inheritance in Man, OMIM™. 2002. Johns Hopkins University, Baltimore, MD. MIM Number: 605086. World Wide Web URL: <http://www.ncbi.nlm.nih.gov/omim/>
5. LocusLink Report (LocusID: 54209). <http://www.ncbi.nlm.nih.gov/LocusLink/>

## CHROMOSOMAL LOCATION

Genetic locus: TREM2 (human) mapping to 6p21.1.

## SOURCE

TREM-2 (H-160) is a rabbit polyclonal antibody raised against amino acids 1-160 mapping at the N-terminus of TREM-2 of human origin.

## PRODUCT

Each vial contains 200 µg IgG in 1.0 ml of PBS with < 0.1% sodium azide and 0.1% gelatin.

## STORAGE

Store at 4° C, \*\*DO NOT FREEZE\*\*. Stable for one year from the date of shipment. Non-hazardous. No MSDS required.

## PROTOCOLS

See our web site at [www.scbt.com](http://www.scbt.com) or our catalog for detailed protocols and support products.

## APPLICATIONS

TREM-2 (H-160) is recommended for detection of TREM-2 isoforms 1-3 of human origin by Western Blotting (starting dilution 1:200, dilution range 1:100-1:1000), immunoprecipitation [1-2 µg per 100-500 µg of total protein (1 ml of cell lysate)], immunofluorescence (starting dilution 1:50, dilution range 1:50-1:500) and solid phase ELISA (starting dilution 1:30, dilution range 1:30-1:3000).

Suitable for use as control antibody for TREM-2 siRNA (h): sc-43001, TREM-2 shRNA Plasmid (h): sc-43001-SH and TREM-2 shRNA (h) Lentiviral Particles: sc-43001-V.

Molecular Weight of glycosylated TREM-2: 40 kDa.


Molecular Weight of deglycosylated TREM-2: 26 kDa.

Positive Controls: HeLa whole cell lysate: sc-2200, Jurkat whole cell lysate: sc-2204 or LNCaP cell lysate: sc-2231.


## RECOMMENDED SECONDARY REAGENTS

To ensure optimal results, the following support (secondary) reagents are recommended: 1) Western blotting: use goat anti-rabbit IgG-HRP: sc-2004 (dilution range: 1:2000-1:100,000) or Cruz Marker™ compatible goat anti-rabbit IgG-HRP: sc-2030 (dilution range: 1:2000-1:5000), Cruz Marker™ Molecular Weight Standards: sc-2035, TBS Blotto A Blocking Reagent: sc-2333 and Western Blotting Luminol Reagent: sc-2048. 2) Immunoprecipitation: use Protein A/G PLUS-Agarose: sc-2003 (0.5 ml agarose/2.0 ml). 3) Immunofluorescence: use goat anti-rabbit IgG-FITC: sc-2012 (dilution range: 1:100-1:400) or goat anti-rabbit IgG-TR: sc-2780 (dilution range: 1:100-1:400) with UltraCruz™ Mounting Medium: sc-24941.

## DATA



TREM-2 (H-160): sc-48764. Western blot analysis of TREM-2 expression in Jurkat (A), LNCaP (B), HeLa (C) and THP-1 (D) whole cell lysates and human liver tissue extract (E).



TREM-2 (H-160): sc-48764. Western blot analysis of TREM-2 expression in non transfected (A) and human TREM-2 transfected (B) HEK293T whole cell lysates.

## RESEARCH USE

For research use only, not for use in diagnostic procedures.

**MONOS**  
Satisfaction  
Guaranteed

Try **TREM-2 (B-3): sc-373828**, our highly recommended monoclonal alternative to TREM-2 (H-160). Also, for AC, HRP, FITC, PE, Alexa Fluor® 488 and Alexa Fluor® 647 conjugates, see **TREM-2 (B-3): sc-373828**.