

POMT2 (N-18): sc-48919

BACKGROUND

O-mannosylation is an essential protein modification in eukaryotes that is initiated by an evolutionarily conserved family of protein O-mannosyltransferases. POMT2 encodes an integral membrane protein which localizes to the endoplasmic reticulum (ER) and shares significant sequence similarity with a family of protein O-mannosyltransferases of *S. cerevisiae*. The deduced 750 amino acid protein has a seven transmembrane helical structure with a central hydrophilic domain surrounded by five N-terminal and two C-terminal transmembrane regions. Like other known members of its family, POMT2 lacks a characteristic ER-targeting or -retention signal and contains five N-glycosylation sites. POMT2 shares 36% sequence identity with human POMT1 and RNA dot blot analysis reveals highest expression of mouse POMT2 in testis.

REFERENCES

1. Willer, T., et al. 2002. Characterization of POMT2, a novel member of the PMT protein O-mannosyltransferase family specifically localized to the acrosome of mammalian spermatids. *Glycobiology* 12: 771-783.
2. Akasaka-Manya, K., et al. 2004. Mutations of the POMT1 gene found in patients with Walker-Warburg syndrome lead to a defect of protein O-mannosylation. *Biochem. Biophys. Res. Commun.* 325: 75-79.
3. Ichimiya, T., et al. 2004. The twisted abdomen phenotype of *Drosophila* POMT1 and POMT2 mutants coincides with their heterophilic protein O-mannosyltransferase activity. *J. Biol. Chem.* 279: 42638-42647.
4. Manya, H., et al. 2004. Demonstration of mammalian protein O-mannosyltransferase activity: coexpression of POMT1 and POMT2 required for enzymatic activity. *Proc. Natl. Acad. Sci. USA* 101: 500-505.
5. van Reeuwijk, J., et al. 2005. POMT2 mutations cause α -dystroglycan hypoglycosylation and Walker-Warburg syndrome. *J. Med. Genet.* 42: 907-912.
6. Manya, H., et al. 2006. Molecular cloning and characterization of rat POMT1 and POMT2. *Glycobiology* 16: 863-873.
7. Mercuri, E., et al. 2006. POMT2 mutation in a patient with "MEB-like" phenotype. *Neuromuscul. Disord.* 16: 446-448.

CHROMOSOMAL LOCATION

Genetic locus: POMT2 (human) mapping to 14q24.3; Pomt2 (mouse) mapping to 12 D2.

SOURCE

POMT2 (N-18) is an affinity purified goat polyclonal antibody raised against a peptide mapping within an internal region of POMT2 of human origin.

PRODUCT

Each vial contains 200 μ g IgG in 1.0 ml of PBS with < 0.1% sodium azide and 0.1% gelatin.

Blocking peptide available for competition studies, sc-48919 P, (100 μ g peptide in 0.5 ml PBS containing < 0.1% sodium azide and 0.2% BSA).

APPLICATIONS

POMT2 (N-18) is recommended for detection of POMT2 of mouse, rat and human origin by Western Blotting (starting dilution 1:200, dilution range 1:100-1:1000), immunoprecipitation [1-2 μ g per 100-500 μ g of total protein (1 ml of cell lysate)], immunofluorescence (starting dilution 1:50, dilution range 1:50-1:500) and solid phase ELISA (starting dilution 1:30, dilution range 1:30-1:3000).


POMT2 (N-18) is also recommended for detection of POMT2 in additional species, including equine, canine, bovine, porcine and avian.

Suitable for use as control antibody for POMT2 siRNA (h): sc-61381, POMT2 siRNA (m): sc-61382, POMT2 shRNA Plasmid (h): sc-61381-SH, POMT2 shRNA Plasmid (m): sc-61382-SH, POMT2 shRNA (h) Lentiviral Particles: sc-61381-V and POMT2 shRNA (m) Lentiviral Particles: sc-61382-V.


Molecular Weight of POMT2: 87 kDa.

Positive Controls: POMT2 (h): 293T Lysate: sc-114414 or MCF7 whole cell lysate: sc-2206.

DATA



POMT2 (N-18): sc-48919. Western blot analysis of POMT2 expression in MCF7 whole cell lysate.



POMT2 (N-18): sc-48919. Western blot analysis of POMT2 expression in non-transfected: sc-117752 (A) and human POMT2 transfected: sc-114414 (B) 293T whole cell lysates.

SELECT PRODUCT CITATIONS

1. Kim, H.W., et al. 2009. Ischemic preconditioning augments survival of stem cells via miR-210 expression by targeting caspase-8-associated protein 2. *J. Biol. Chem.* 284: 33161-33168.

STORAGE

Store at 4° C, ****DO NOT FREEZE****. Stable for one year from the date of shipment. Non-hazardous. No MSDS required.

RESEARCH USE

For research use only, not for use in diagnostic procedures.



Try **POMT2 (G-3): sc-393487**, our highly recommended monoclonal alternative to POMT2 (N-18).