

RasGRP3 (R-17): sc-48926

BACKGROUND

Members of the Ras subfamily of GTPases function in signal transduction as GTP/GDP-modulated switches that rotate between inactive GDP- and active GTP-bound states. Guanine nucleotide exchange factors (GEFs), such as RasGRP3 (GRP3), act as Ras activators by promoting retrieval of GTP to maintain the active GTP-bound state and are the fundamental link between cell surface receptors and Ras activation. Highest levels of RasGRP3 expression are observed in heart, brain, lung and kidney tissues, and intermediate expression is observed in liver, skeletal muscle, pancreas, spleen, testis and ovary tissues. RasGRP3, which shares significant sequence identity with the calcium- and diacylglycerol-activated GEFs, activates Ras and Rap 1 and promotes activation of ELK1 in prostate cancer cell lines.

REFERENCES

- Nagase, T., et al. 1999. Prediction of the coding sequences of unidentified human genes. XII. The complete sequences of 100 new cDNA clones from brain which code for large proteins *in vitro*. DNA Res. 5: 355-364.
- Rebhun, J.F., et al. 2000. Identification of guanine nucleotide exchange factors (GEFs) for the Rap 1 GTPase. Regulation of MR-GEF by M-Ras-GTP interaction. J. Biol. Chem. 275: 34901-34908.
- Aiba, Y., et al. 2004. Activation of RasGRP3 by phosphorylation of Thr 133 is required for B cell receptor-mediated Ras activation. Proc. Natl. Acad. Sci. USA 101: 16612-16617.
- Roberts, D.M., et al. 2004. A vascular gene trap screen defines RasGRP3 as an angiogenesis-regulated gene required for the endothelial response to phorbol esters. Mol. Cell. Biol. 24: 10515-10528.
- Braun, D.C., et al. 2005. Role of phorbol ester localization in determining protein kinase C or RasGRP3 translocation: real-time analysis using fluorescent ligands and proteins. Mol. Cancer Ther. 4: 141-150.
- Coughlin, J.J., et al. 2005. RasGRP1 and RasGRP3 regulate B cell proliferation by facilitating B cell receptor-Ras signaling. J. Immunol. 175: 7179-7184.
- Ozaki, N., et al. 2005. RasGRP3 mediates phorbol ester-induced, protein kinase C-independent exocytosis. Biochem. Biophys. Res. Commun. 329: 765-771.
- Regier, D.S., et al. 2005. Diacylglycerol kinase α regulates Ras guanyl-releasing protein 3 and inhibits Rap 1 signaling. Proc. Natl. Acad. Sci. USA 102: 7595-7600.
- Zheng, Y., et al. 2005. Phosphorylation of RasGRP3 on Threonine 133 provides a mechanistic link between PKC and Ras signaling systems in B cells. Blood 105: 3648-3654.

CHROMOSOMAL LOCATION

Genetic locus: RASGRP3 (human) mapping to 2p22.3; Rasgrp3 (mouse) mapping to 17 E2.

SOURCE

RasGRP3 (R-17) is an affinity purified goat polyclonal antibody raised against a peptide mapping near the N-terminus of RasGRP3 of human origin.

PRODUCT

Each vial contains 200 μ g IgG in 1.0 ml of PBS with < 0.1% sodium azide and 0.1% gelatin.

Blocking peptide available for competition studies, sc-48926 P, (100 μ g peptide in 0.5 ml PBS containing < 0.1% sodium azide and 0.2% BSA).

APPLICATIONS

RasGRP3 (R-17) is recommended for detection of RasGRP3 of mouse, rat and human origin by Western Blotting (starting dilution 1:200, dilution range 1:100-1:1000), immunofluorescence (starting dilution 1:50, dilution range 1:50-1:500) and solid phase ELISA (starting dilution 1:30, dilution range 1:30-1:3000).

RasGRP3 (R-17) is also recommended for detection of RasGRP3 in additional species, including equine, canine, bovine, porcine and avian.

Suitable for use as control antibody for RasGRP3 siRNA (h): sc-61444, RasGRP3 siRNA (m): sc-61445, RasGRP3 shRNA Plasmid (h): sc-61444-SH, RasGRP3 shRNA Plasmid (m): sc-61445-SH, RasGRP3 shRNA (h) Lentiviral Particles: sc-61444-V and RasGRP3 shRNA (m) Lentiviral Particles: sc-61445-V.

Molecular Weight of RasGRP3: 80 kDa.

Positive Controls: Ramos cell lysate: sc-2216.

RECOMMENDED SECONDARY REAGENTS

To ensure optimal results, the following support (secondary) reagents are recommended: 1) Western Blotting: use donkey anti-goat IgG-HRP: sc-2020 (dilution range: 1:2000-1:100,000) or Cruz Marker™ compatible donkey anti-goat IgG-HRP: sc-2033 (dilution range: 1:2000-1:5000), Cruz Marker™ Molecular Weight Standards: sc-2035, TBS Blotto A Blocking Reagent: sc-2333 and Western Blotting Luminol Reagent: sc-2048. 2) Immunofluorescence: use donkey anti-goat IgG-FITC: sc-2024 (dilution range: 1:100-1:400) or donkey anti-goat IgG-TR: sc-2783 (dilution range: 1:100-1:400) with UltraCruz™ Mounting Medium: sc-24941.

STORAGE

Store at 4° C, ****DO NOT FREEZE****. Stable for one year from the date of shipment. Non-hazardous. No MSDS required.

RESEARCH USE

For research use only, not for use in diagnostic procedures.

PROTOCOLS

See our web site at www.scbt.com or our catalog for detailed protocols and support products.