



CYP2R1 (T-12): sc-48987

BACKGROUND

P450 enzymes constitute a family of monooxygenase enzymes that are involved in the metabolism of a wide array of endogenous and xenobiotic compounds. There are currently 57 known active cytochrome P450 (CYP) genes and 58 known pseudogenes present in the human genome. CYP2R1 is a 501 amino acid protein which belongs to the CYP2 family of cytochrome P450 proteins. These proteins are usually involved in the metabolism of foreign compounds. CYP2R1 acts as a major vitamin D 25-hydroxylase. A homozygous mutation in the CYP2R1 gene eliminates the vitamin D3 25-hydroxylase activity of the CYP2R1 protein, causing 25-hydroxyvitamin D3 deficiency. Manifestations of 25-hydroxyvitamin D3 deficiency include abnormally low plasma levels of 25-hydroxyvitamin D3 and various symptoms of vitamin D deficiency, including skeletal abnormalities, hypophosphatemia and hypocalcemia.

REFERENCES

1. Ponchon, G., Kennan, A.L. and DeLuca, H.F. 1969. "Activation" of vitamin D by the liver. *J. Clin. Invest.* 48: 2032-2037.
2. Online Mendelian Inheritance in Man, OMIM™. 2002. Johns Hopkins University, Baltimore, MD. MIM Number: 608713. World Wide Web URL: <http://www.ncbi.nlm.nih.gov/omim/>
3. Cheng, J.B., Levine, M.A., Bell, N.H., Mangelsdorf, D.J. and Russell, D.W. 2004. Genetic evidence that the human CYP2R1 enzyme is a key vitamin D 25-hydroxylase. *Proc. Natl. Acad. Sci. USA* 101: 7711-7715.
4. Prosser, D.E. and Jones, G. 2004. Enzymes involved in the activation and inactivation of vitamin D. *Trends Biochem. Sci.* 29: 664-673.
5. Shinkyo, R., Sakaki, T., Kamakura, M., Ohta, M. and Inouye, K. 2004. Metabolism of vitamin D by human microsomal CYP2R1. *Biochem. Biophys. Res. Commun.* 324: 451-457.
6. Choudhary, D., Jansson, I., Stoilov, I., Sarfarazi, M. and Schenkman, J.B. 2005. Expression patterns of mouse and human CYP orthologs (families 1-4) during development and in different adult tissues. *Arch. Biochem. Biophys.* 436: 50-61.
7. Ingelman-Sundberg, M. 2005. The human genome project and novel aspects of cytochrome P450 research. *Toxicol. Appl. Pharmacol.* 207: 52-56.
8. Karlgren, M., Miura, S. and Ingelman-Sundberg, M. 2005. Novel extra-hepatic cytochrome P450s. *Toxicol. Appl. Pharmacol.* 207: 57-61.

CHROMOSOMAL LOCATION

Genetic locus: CYP2R1 (human) mapping to 11p15.2; Cyp2r1 (mouse) mapping to 7 F1.

SOURCE

CYP2R1 (T-12) is an affinity purified goat polyclonal antibody raised against a peptide mapping near the C-terminus of CYP2R1 of human origin.

STORAGE

Store at 4° C, ****DO NOT FREEZE****. Stable for one year from the date of shipment. Non-hazardous. No MSDS required.

PRODUCT

Each vial contains 200 µg IgG in 1.0 ml of PBS with < 0.1% sodium azide and 0.1% gelatin.

Blocking peptide available for competition studies, sc-48987 P, (100 µg peptide in 0.5 ml PBS containing < 0.1% sodium azide and 0.2% BSA).

APPLICATIONS

CYP2R1 (T-12) is recommended for detection of CYP2R1 of mouse, rat and human origin by Western Blotting (starting dilution 1:200, dilution range 1:100-1:1000), immunofluorescence (starting dilution 1:50, dilution range 1:50-1:500) and solid phase ELISA (starting dilution 1:30, dilution range 1:30-1:3000).

Suitable for use as control antibody for CYP2R1 siRNA (h): sc-60481 and CYP2R1 siRNA (m): sc-60482.

Molecular Weight of CYP2R1: 57 kDa.

RECOMMENDED SECONDARY REAGENTS

To ensure optimal results, the following support (secondary) reagents are recommended: 1) Western Blotting: use donkey anti-goat IgG-HRP: sc-2020 (dilution range: 1:2000-1:100,000) or Cruz Marker™ compatible donkey anti-goat IgG-HRP: sc-2033 (dilution range: 1:2000-1:5000), Cruz Marker™ Molecular Weight Standards: sc-2035, TBS Blotto A Blocking Reagent: sc-2333 and Western Blotting Luminol Reagent: sc-2048. 2) Immunofluorescence: use donkey anti-goat IgG-FITC: sc-2024 (dilution range: 1:100-1:400) or donkey anti-goat IgG-TR: sc-2783 (dilution range: 1:100-1:400) with UltraCruz™ Mounting Medium: sc-24941.

RESEARCH USE

For research use only, not for use in diagnostic procedures.

PROTOCOLS

See our web site at www.scbt.com or our catalog for detailed protocols and support products.