

MCAD (F-15): sc-49046

BACKGROUND

Acyl-CoA dehydrogenase is a family of enzymes that localize to the mitochondrion and target acyl chain lengths of 4 to 16 by use of the mitochondrial fatty acid β -oxidation pathway. In mammalian tissue, many straight-chain acyl-CoA dehydrogenases possess different substrate specificities. In rare cases, irregularities in medium-chain acyl-CoA dehydrogenase can cause fasting hypoglycemia, hepatic dysfunction and encephalopathy, often resulting in death in infancy. MCAD, also designated acyl-CoA dehydrogenase, medium-chain (ACADM) and MCADH, is a homotetramer. The MCAD gene encodes a 421 amino acid protein with characteristics of mitochondrial protein transit peptides. The protein shows 88% sequence identity with MCAD of porcine origin. Medium-chain acyl-CoA dehydrogenase catalyzes the initial reaction in the β -oxidation of C4 to C12 straight-chain acyl-CoAs.

CHROMOSOMAL LOCATION

Genetic locus: ACADM (human) mapping to 1p31.1; Acadm (mouse) mapping to 3 H3.

SOURCE

MCAD (F-15) is an affinity purified goat polyclonal antibody raised against a peptide mapping near the N-terminus of MCAD of human origin.

PRODUCT

Each vial contains 200 μ g IgG in 1.0 ml of PBS with < 0.1% sodium azide and 0.1% gelatin.

Blocking peptide available for competition studies, sc-49046 P, (100 μ g peptide in 0.5 ml PBS containing < 0.1% sodium azide and 0.2% BSA).

STORAGE

Store at 4° C, ****DO NOT FREEZE****. Stable for one year from the date of shipment. Non-hazardous. No MSDS required.

APPLICATIONS

MCAD (F-15) is recommended for detection of MCAD of mouse, rat and human origin by Western Blotting (starting dilution 1:200, dilution range 1:100-1:1000), immunoprecipitation [1-2 μ g per 100-500 μ g of total protein (1 ml of cell lysate)], immunofluorescence (starting dilution 1:50, dilution range 1:50-1:500) and solid phase ELISA (starting dilution 1:30, dilution range 1:30-1:3000).

MCAD (F-15) is also recommended for detection of MCAD in additional species, including equine, canine, bovine and porcine.

Suitable for use as control antibody for MCAD siRNA (h): sc-60996, MCAD siRNA (m): sc-60997, MCAD shRNA Plasmid (h): sc-60996-SH, MCAD shRNA Plasmid (m): sc-60997-SH, MCAD shRNA (h) Lentiviral Particles: sc-60996-V and MCAD shRNA (m) Lentiviral Particles: sc-60997-V.


Molecular Weight of MCAD: 45 kDa.

Positive Controls: MCAD (h2): 293T Lysate: sc-170633, Hep G2 cell lysate: sc-2227 or mouse heart extract: sc-2254.


RECOMMENDED SECONDARY REAGENTS

To ensure optimal results, the following support (secondary) reagents are recommended: 1) Western Blotting: use donkey anti-goat IgG-HRP: sc-2020 (dilution range: 1:2000-1:100,000) or Cruz Marker™ compatible donkey anti-goat IgG-HRP: sc-2033 (dilution range: 1:2000-1:5000), Cruz Marker™ Molecular Weight Standards: sc-2035, TBS Blotto A Blocking Reagent: sc-2333 and Western Blotting Luminol Reagent: sc-2048. 2) Immunoprecipitation: use Protein A/G PLUS-Agarose: sc-2003 (0.5 ml agarose/2.0 ml). 3) Immunofluorescence: use donkey anti-goat IgG-FITC: sc-2024 (dilution range: 1:100-1:400) or donkey anti-goat IgG-TR: sc-2783 (dilution range: 1:100-1:400) with UltraCruz™ Mounting Medium: sc-24941.

DATA



MCAD (F-15): sc-49046. Western blot analysis of MCAD expression in non-transfected 293T: sc-117752 (A), human MCAD transfected 293T: sc-170633 (B) and Hep G2 (C) whole cell lysates.



MCAD (F-15): sc-49046. Immunofluorescence staining of methanol-fixed HeLa cells showing mitochondrial localization.

RESEARCH USE

For research use only, not for use in diagnostic procedures.

PROTOCOLS

See our web site at www.scbt.com or our catalog for detailed protocols and support products.



Try **MCAD (E-8): sc-271931** or **MCAD (E-8): sc-271931**, our highly recommended monoclonal alternatives to MCAD (F-15).