

SNX7 (C-17): sc-49134

BACKGROUND

Sorting nexin (SNX) proteins are members of a large family of hydrophilic PX (phospholipid-binding motif) domain-containing proteins that interact with a variety of receptor types. SNXs are widely expressed, although the tissue distribution of each SNX mRNA varies. The ability of SNXs to bind specific phospholipids, as well as their tendency to form protein-protein complexes, suggests a role for these proteins in cellular membrane trafficking and protein sorting. SNXs may also function specifically in pro-degradative sorting, internalization, endosomal recycling or simply in endosomal sorting. SNXs partially associate with cellular membranes, despite their hydrophilic nature. SNX7 is unique in that it does not have a coiled coil region like some of the SNX family members. Mutations in the SNX7 gene have not been shown to cause any diseases.

REFERENCES

1. Worby, C.A. and Dixon, J.E. 2002. Sorting out the cellular functions of sorting nexins. *Nat. Rev. Mol. Cell Biol.* 3: 919-931.
2. Orlacchio, A., Kawarai, T., Gaudiello, F., St. George-Hyslop, P.H., Floris, R. and Bernardi, G. 2005. New locus for hereditary spastic paraplegia maps to chromosome 1p31.1-1p21.1. *Ann. Neurol.* 58: 423-429.
3. Carlton, J.G. and Cullen, P.J. 2005. Sorting nexins. *Curr. Biol.* 15: 819-820.
4. Carlton, J., Bujny, M., Rutherford, A. and Cullen, P. 2005. Sorting nexins: unifying trends and new perspectives. *Traffic* 6: 75-82.
5. Jacques, C., Baris, O., Prunier-Mirebeau, D., Savagner, F., Rodien, P., Rohmer, V., Franc, B., Guyetant, S., Malthiery, Y. and Reynier, P. 2005. Two-step differential expression analysis reveals a new set of genes involved in thyroid oncocytic tumors. *J. Clin. Endocrinol. Metab.* 90: 2314-2320.
6. Kerr, M.C., Lindsay, M.R., Luetterforst, R., Hamilton, N., Simpson, F., Parton, R.G., Gleeson, P.A. and Teasdale, R.D. 2006. Visualisation of macropinosome maturation by the recruitment of sorting nexins. *J. Cell Sci.* 119: 3967-3980.

CHROMOSOMAL LOCATION

Genetic locus: SNX7 (human) mapping to 1p21.3; Snx7 (mouse) mapping to 3 G1.

SOURCE

SNX7 (C-17) is an affinity purified goat polyclonal antibody raised against a peptide mapping at the C-terminus of SNX7 of human origin.

PRODUCT

Each vial contains 200 µg IgG in 1.0 ml of PBS with < 0.1% sodium azide and 0.1% gelatin.

Blocking peptide available for competition studies, sc-49134 P, (100 µg peptide in 0.5 ml PBS containing < 0.1% sodium azide and 0.2% BSA).

STORAGE

Store at 4° C, ****DO NOT FREEZE****. Stable for one year from the date of shipment. Non-hazardous. No MSDS required.

APPLICATIONS

SNX7 (C-17) is recommended for detection of SNX7 of mouse, rat and human origin by Western Blotting (starting dilution 1:200, dilution range 1:100-1:1000), immunofluorescence (starting dilution 1:50, dilution range 1:50-1:500), immunohistochemistry (including paraffin-embedded sections) (starting dilution 1:50, dilution range 1:50-1:500) and solid phase ELISA (starting dilution 1:30, dilution range 1:30-1:3000).

SNX7 (C-17) is also recommended for detection of SNX7 in additional species, including equine, bovine and porcine.

Suitable for use as control antibody for SNX7 siRNA (h): sc-61593, SNX7 siRNA (m): sc-61594, SNX7 shRNA Plasmid (h): sc-61593-SH, SNX7 shRNA Plasmid (m): sc-61594-SH, SNX7 shRNA (h) Lentiviral Particles: sc-61593-V and SNX7 shRNA (m) Lentiviral Particles: sc-61594-V.


Molecular Weight of SNX7 isoforms 1-3: 45/39/52 kDa.

Positive Controls: HeLa whole cell lysate: sc-2200 or MCF7 whole cell lysate: sc-2206.

RECOMMENDED SECONDARY REAGENTS

To ensure optimal results, the following support (secondary) reagents are recommended: 1) Western Blotting: use donkey anti-goat IgG-HRP: sc-2020 (dilution range: 1:2000-1:100,000) or Cruz Marker™ compatible donkey anti-goat IgG-HRP: sc-2033 (dilution range: 1:2000-1:5000), Cruz Marker™ Molecular Weight Standards: sc-2035, TBS Blotto A Blocking Reagent: sc-2333 and Western Blotting Luminol Reagent: sc-2048. 2) Immunofluorescence: use donkey anti-goat IgG-FITC: sc-2024 (dilution range: 1:100-1:400) or donkey anti-goat IgG-TR: sc-2783 (dilution range: 1:100-1:400) with UltraCruz™ Mounting Medium: sc-24941. 3) Immunohistochemistry: use ImmunoCruz™: sc-2053 or ABC: sc-2023 goat IgG Staining Systems.

DATA



SNX7 (C-17): sc-49134. Immunoperoxidase staining of formalin fixed, paraffin-embedded human heart muscle tissue showing cytoplasmic staining of myocytes.

RESEARCH USE

For research use only, not for use in diagnostic procedures.

PROTOCOLS

See our web site at www.scbt.com or our catalog for detailed protocols and support products.