

nephrocystin-4 (T-17): sc-49247

BACKGROUND

The nephrocystin proteins comprise a family of five enzymes that commonly interact with p130Cas, proline-rich tyrosine kinases, calmodulin and tensin, indicating that these proteins may participate in a common signaling pathway. nephrocystin-4 is a 1,250 amino acid protein that interacts with signaling molecules involved in cell adhesion and organization of the actin cytoskeleton, such as Pyk2, tensin and filamins. nephrocystin-4 co-localizes with PKD-2 in the transition zones of ciliated sensory endings of dendrites, and together they play an important role in facilitating ciliary sensory signal transduction. Mutations in the nephrocystin-4 gene contribute to the disease nephronophthisis, an autosomal-recessive cystic kidney disease. Clinical features of familial juvenile nephronophthisis include anemia, polyuria, polydipsia, isostenuria and death.

REFERENCES

1. Mollet, G., et al. 2002. The gene mutated in juvenile nephronophthisis type 4 encodes a novel protein that interacts with nephrocystin. *Nat. Genet.* 32: 300-305.
2. Online Mendelian Inheritance in Man, OMIM™. 2002. Johns Hopkins University, Baltimore, MD. MIM Number: 607215. World Wide Web URL: <http://www.ncbi.nlm.nih.gov/omim/>
3. Olbrich, H., et al. 2003. Mutations in a novel gene, NPHP3, cause adolescent nephronophthisis, tapeto-retinal degeneration and hepatic fibrosis. *Nat. Genet.* 34: 455-459.
4. Hoefele, J., et al. 2004. Clinical and histological presentation of three siblings with mutations in the NPHP4 gene. *Am. J. Kidney Dis.* 43: 358-364.
5. Jauregui, A.R., et al. 2005. Functional characterization of the *C. elegans* nephrocystins NPHP-1 and NPHP-4 and their role in cilia and male sensory behaviors. *Exp. Cell Res.* 305: 333-342.
6. Mollet, G., et al. 2005. Characterization of the nephrocystin/Nephrocystin-4 complex and subcellular localization of Nephrocystin-4 to primary cilia and centrosomes. *Hum. Mol. Genet.* 14: 645-656.
7. Roepman, R., et al. 2005. Interaction of Nephrocystin-4 and RPGRIP1 is disrupted by nephronophthisis or Leber congenital amaurosis-associated mutations. *Proc. Natl. Acad. Sci. USA* 102: 18520-18525.
8. Winkelbauer, M.E., et al. 2005. The *C. elegans* homologs of Nephrocystin-1 and Nephrocystin-4 are cilia transition zone proteins involved in chemosensory perception. *J. Cell Sci.* 118: 5575-5587.
9. Wolf, M.T., et al. 2005. Expression and phenotype analysis of the Nephrocystin-1 and Nephrocystin-4 homologs in *Caenorhabditis elegans*. *J. Am. Soc. Nephrol.* 16: 676-687.

CHROMOSOMAL LOCATION

Genetic locus: NPHP4 (human) mapping to 1p36.31; Nphp4 (mouse) mapping to 4 E2.

STORAGE

Store at 4° C, **DO NOT FREEZE**. Stable for one year from the date of shipment. Non-hazardous. No MSDS required.

SOURCE

Nephrocystin-4 (T-17) is an affinity purified goat polyclonal antibody raised against a peptide mapping near the C-terminus of Nephrocystin-4 of human origin.

PRODUCT

Each vial contains 200 µg IgG in 1.0 ml of PBS with < 0.1% sodium azide and 0.1% gelatin.

Blocking peptide available for competition studies, sc-49247 P, (100 µg peptide in 0.5 ml PBS containing < 0.1% sodium azide and 0.2% BSA).

APPLICATIONS

nephrocystin-4 (T-17) is recommended for detection of nephrocystin-4 of mouse, rat and human origin by Western Blotting (starting dilution 1:200, dilution range 1:100-1:1000), immunofluorescence (starting dilution 1:50, dilution range 1:50-1:500) and solid phase ELISA (starting dilution 1:30, dilution range 1:30-1:3000).

nephrocystin-4 (T-17) is also recommended for detection of nephrocystin-4 in additional species, including equine.

Suitable for use as control antibody for nephrocystin-4 siRNA (h): sc-61182, nephrocystin-4 siRNA (m): sc-61183, nephrocystin-4 shRNA Plasmid (h): sc-61182-SH, nephrocystin-4 shRNA Plasmid (m): sc-61183-SH, nephrocystin-4 shRNA (h) Lentiviral Particles: sc-61182-V and nephrocystin-4 shRNA (m) Lentiviral Particles: sc-61183-V.

Molecular Weight of nephrocystin-4: 175 kDa.

RECOMMENDED SECONDARY REAGENTS

To ensure optimal results, the following support (secondary) reagents are recommended: 1) Western Blotting: use donkey anti-goat IgG-HRP: sc-2020 (dilution range: 1:2000-1:100,000) or Cruz Marker™ compatible donkey anti-goat IgG-HRP: sc-2033 (dilution range: 1:2000-1:5000), Cruz Marker™ Molecular Weight Standards: sc-2035, TBS Blotto A Blocking Reagent: sc-2333 and Western Blotting Luminol Reagent: sc-2048. 2) Immunofluorescence: use donkey anti-goat IgG-FITC: sc-2024 (dilution range: 1:100-1:400) or donkey anti-goat IgG-TR: sc-2783 (dilution range: 1:100-1:400) with UltraCruz™ Mounting Medium: sc-24941.

RESEARCH USE

For research use only, not for use in diagnostic procedures.

PROTOCOLS

See our web site at www.scbt.com or our catalog for detailed protocols and support products.