

AXUD1 (D-15): sc-49280

BACKGROUND

Axin, an important regulator of β -catenin, is frequently mutated in human hepatocellular carcinomas (HCCs). Transduction of the wildtype Axin gene (AXIN1) induces apoptosis in colon cancer cells as well as in HCC cells. Axin-1 upregulated gene 1 protein (AXUD1), also designated TGF β -induced apoptosis protein 3 (TAIP-3) or URAX1, is ubiquitously expressed, but highest levels are detected in placenta, lung, skeletal muscle, leukocytes and pancreas. AXUD1 has increased expression in response to exogenously expressed Axin-1 and localizes primarily to the nucleus. It is downregulated in kidney, lung, liver and colon cancers compared with corresponding unaffected tissues, which implicates that AXUD1 may function as a tumor-suppressor in these organs. AXUD1 may also be involved in apoptosis.

REFERENCES

- Hart, M.J., de los Santos, R., Albert, I.N., Rubinfeld, B. and Polakis, P. 1998. Downregulation of β -catenin by human Axin and its association with the APC tumor suppressor, β -catenin and GSK3 β . *Curr. Biol.* 8: 573-581.
- Satoh, S., Daigo, Y., Furukawa, Y., Kato, T., Miwa, N., Nishiwaki, T., Kawasoe, T., Ishiguro, H., Fujita, M., Tokino, T., Sasaki, Y., Imaoka, S., Murata, M., Shimano, T., Yamaoka, Y. and Nakamura, Y. 2000. AXIN1 mutations in hepatocellular carcinomas, and growth suppression in cancer cells by virus-mediated transfer of AXIN1. *Nat. Genet.* 24: 245-250.
- Ishiguro, H., Tsunoda, T., Tanaka, T., Fujii, Y., Nakamura, Y. and Furukawa, Y. 2001. Identification of AXUD1, a novel human gene induced by AXIN1 and its reduced expression in human carcinomas of the lung, liver, colon and kidney. *Oncogene* 20: 5062-5066.
- Online Mendelian Inheritance in Man, OMIM™. 2002. Johns Hopkins University, Baltimore, MD. MIM Number: 606458. World Wide Web URL: <http://www.ncbi.nlm.nih.gov/omim/>
- Yi, X., Bekeredian, R., DeFilippis, N.J., Siddiquee, Z., Fernandez, E. and Shohet, R.V. 2006. Transcriptional analysis of doxorubicin-induced cardiotoxicity. *Am. J. Physiol. Heart Circ. Physiol.* 290: H1098-H1102.

CHROMOSOMAL LOCATION

Genetic locus: CSRNP1 (human) mapping to 3p22.2; Csrnp1 (mouse) mapping to 9 F4.

SOURCE

AXUD1 (D-15) is an affinity purified goat polyclonal antibody raised against a peptide mapping near the N-terminus of AXUD1 of human origin.

PRODUCT

Each vial contains 200 μ g IgG in 1.0 ml of PBS with < 0.1% sodium azide and 0.1% gelatin.

Blocking peptide available for competition studies, sc-49280 P, (100 μ g peptide in 0.5 ml PBS containing < 0.1% sodium azide and 0.2% BSA).

RESEARCH USE

For research use only, not for use in diagnostic procedures.

APPLICATIONS

AXUD1 (D-15) is recommended for detection of Axin-1 up-regulated gene 1 protein of mouse, rat and human origin by Western Blotting (starting dilution 1:200, dilution range 1:100-1:1000), immunofluorescence (starting dilution 1:50, dilution range 1:50-1:500) and solid phase ELISA (starting dilution 1:30, dilution range 1:30-1:3000).

AXUD1 (D-15) is also recommended for detection of Axin-1 up-regulated gene 1 protein in additional species, including equine, canine, bovine and porcine.

Suitable for use as control antibody for AXUD1 siRNA (h): sc-60237, AXUD1 siRNA (m): sc-60238, AXUD1 shRNA Plasmid (h): sc-60237-SH, AXUD1 shRNA Plasmid (m): sc-60238-SH, AXUD1 shRNA (h) Lentiviral Particles: sc-60237-V and AXUD1 shRNA (m) Lentiviral Particles: sc-60238-V.

Molecular Weight of AXUD1: 64 kDa.

Positive Controls: HeLa whole cell lysate: sc-2200.

RECOMMENDED SECONDARY REAGENTS

To ensure optimal results, the following support (secondary) reagents are recommended: 1) Western Blotting: use donkey anti-goat IgG-HRP: sc-2020 (dilution range: 1:2000-1:100,000) or Cruz Marker™ compatible donkey anti-goat IgG-HRP: sc-2033 (dilution range: 1:2000-1:5000), Cruz Marker™ Molecular Weight Standards: sc-2035, TBS Blotto A Blocking Reagent: sc-2333 and Western Blotting Luminol Reagent: sc-2048. 2) Immunofluorescence: use donkey anti-goat IgG-FITC: sc-2024 (dilution range: 1:100-1:400) or donkey anti-goat IgG-TR: sc-2783 (dilution range: 1:100-1:400) with UltraCruz™ Mounting Medium: sc-24941.

STORAGE

Store at 4° C, **DO NOT FREEZE**. Stable for one year from the date of shipment. Non-hazardous. No MSDS required.

PROTOCOLS

See our web site at www.scbt.com or our catalog for detailed protocols and support products.