

Barttin (D-20): sc-49607

BACKGROUND

The BSND gene encodes Barttin, a protein comprised of two putative trans-membrane α helices. Barttin expression is detected in the thin limb and thick ascending limb of the loop of Henle in the kidney, and in the dark cells of the inner ear. The BSND gene is mutated in Bartter syndrome, a genetic disease characterized by hypokalemia, metabolic alkalosis and normal to low blood pressure, which occurs with sensorineural deafness, irreversible hearing loss due to cochlear sensorineural or cochlear nerve damage. Barttin acts as an essential β subunit for CLCKNA and CLCKNB chloride channels, with which it co-localizes in basolateral membranes of renal tubules and of potassium-secreting epithelia of the inner ear. Mutations in either CLCKNB or Barttin compromise currents through heteromeric channels that can be stimulated further by mutating a proline-tyrosine (PY) motif on Barttin. Heteromers formed by chloride channels and Barttin are essential for renal salt reabsorption and potassium recycling in the inner ear.

REFERENCES

- Estevez, R., Boettger, T., Stein, V., Birkenhager, R., Otto, E., Hildebrandt, F. and Jentsch, T.J. 2001. Barttin is a Cl^- channel β subunit crucial for renal Cl^- reabsorption and inner ear K^+ secretion. *Nature* 414: 558-561.
- Miyamura, N., Matsumoto, K., Taguchi, T., Tokunaga, H., Nishikawa, T., Nishida, K., Toyonaga, T., Sakakida, M. and Araki, E. 2003. Atypical Bartter syndrome with sensorineural deafness with G47R mutation of the β subunit for ClC-Ka and ClC-Kb chloride channels, Barttin. *J. Clin. Endocrinol. Metab.* 88: 781-786.
- Wolf, K., Meier-Meitingner, M., Bergler, T., Castrop, H., Vitzthum, H., Riegger, G.A., Kurtz, A. and Kramer, B.K. 2003. Parallel downregulation of chloride channel CLCK1 and Barttin mRNA in the thin ascending limb of the rat nephron by furosemide. *Pflugers Arch.* 446: 665-671.
- Liantonio, A., Pusch, M., Picollo, A., Guida, P., De Luca, A., Pierno, S., Fracchiolla, G., Loidice, F., Tortorella, P. and Conte Camerino, D. 2004. Investigations of pharmacologic properties of the renal CLCK1 chloride channel co-expressed with Barttin by the use of 2-(p-Chlorophenoxy) propionic acid derivatives and other structurally unrelated chloride channels blockers. *J. Am. Soc. Nephrol.* 15: 13-20.
- Embark, H.M., Bohmer, C., Palmada, M., Rajamanickam, J., Wyatt, A.W., Wallisch, S., Capasso, G., Waldegger, P., Seyberth, H.W., Waldegger, S. and Lang, F. 2004. Regulation of CLCKA/Barttin by the ubiquitin ligase Nedd4-2 and the serum- and glucocorticoid-dependent kinases. *Kidney Int.* 66: 1918-1925.
- Lang, F., Capasso, G., Schwab, M. and Waldegger, S. 2005. Renal tubular transport and the genetic basis of hypertensive disease. *Clin. Exp. Nephrol.* 9: 91-99.

CHROMOSOMAL LOCATION

Genetic locus: BSND (human) mapping to 1p32.3.

SOURCE

Barttin (D-20) is an affinity purified goat polyclonal antibody raised against a peptide mapping within an internal region of Barttin of human origin.

PRODUCT

Each vial contains 200 μg IgG in 1.0 ml of PBS with < 0.1% sodium azide and 0.1% gelatin.

Blocking peptide available for competition studies, sc-49607 P, (100 μg peptide in 0.5 ml PBS containing < 0.1% sodium azide and 0.2% BSA).

APPLICATIONS

Barttin (D-20) is recommended for detection of Barttin isoform 1 of human origin by Western Blotting (starting dilution 1:200, dilution range 1:100-1:1000), immunofluorescence (starting dilution 1:50, dilution range 1:50-1:500) and solid phase ELISA (starting dilution 1:30, dilution range 1:30-1:3000).

Suitable for use as control antibody for Barttin siRNA (h): sc-60245, Barttin shRNA Plasmid (h): sc-60245-SH and Barttin shRNA (h) Lentiviral Particles: sc-60245-V.

Molecular Weight of Barttin: 35 kDa.

RECOMMENDED SECONDARY REAGENTS

To ensure optimal results, the following support (secondary) reagents are recommended: 1) Western Blotting: use donkey anti-goat IgG-HRP: sc-2020 (dilution range: 1:2000-1:100,000) or Cruz Marker™ compatible donkey anti-goat IgG-HRP: sc-2033 (dilution range: 1:2000-1:5000), Cruz Marker™ Molecular Weight Standards: sc-2035, TBS Blotto A Blocking Reagent: sc-2333 and Western Blotting Luminol Reagent: sc-2048. 2) Immunofluorescence: use donkey anti-goat IgG-FITC: sc-2024 (dilution range: 1:100-1:400) or donkey anti-goat IgG-TR: sc-2783 (dilution range: 1:100-1:400) with UltraCruz™ Mounting Medium: sc-24941.

STORAGE

Store at 4° C, **DO NOT FREEZE**. Stable for one year from the date of shipment. Non-hazardous. No MSDS required.

RESEARCH USE

For research use only, not for use in diagnostic procedures.

PROTOCOLS

See our web site at www.scbt.com or our catalog for detailed protocols and support products.



Try **Barttin (A-1): sc-271867** or **Barttin (A-3): sc-365161**, our highly recommended monoclonal alternatives to Barttin (D-20).