

OAS3 (M-14): sc-49869

BACKGROUND

The 2'- 5'- oligoadenylate synthetase (OAS) family is comprised of four members: OAS1, OAS2, OAS3 and OASL. These proteins are induced by interferons and function to convert ATP into 2'- 5'- linked oligomers of adenosine in the presence of double-stranded RNA and magnesium ions. Copper, iron and zinc ions strongly inhibit the OAS enzymatic activity, while manganese ions can replace magnesium ions as an activator. The OAS family plays a significant role in the inhibition of cellular protein synthesis, apoptosis and growth, and its members are important factors in viral infection resistance. OAS3, also referred to as p100, contains three adjacent OAS1-like domains and maps to the human chromosome 12q24.2.

REFERENCES

- Hovanessian, A.G., Laurent, A.G., Chebath, J., Galabru, J., Robert, N. and Svab, J. 1987. Identification of 69 kDa and 100 kDa forms of 2-5A synthetase in interferon-treated human cells by specific monoclonal antibodies. *EMBO J.* 6: 1273-1280.
- Corrias, M.V., Gribaudo, G., Guarnaccia, F. and Ponzoni, M. 1995. Induction of 2.5 OAS gene expression and activity is not sufficient for IFN- γ -induced neuroblastoma cell differentiation. *Int. J. Cancer* 62: 223-229.
- Hovnanian, A., Rebouillat, D., Mattei, M.G., Levy, E.R., Marie, I., Monaco, A.P. and Hovanessian, A.G. 1998. The human 2'- 5'- oligoadenylate synthetase locus is composed of three distinct genes clustered on chromosome 12q24.2 encoding the 100, 69, and 40 kDa forms. *Genomics* 52: 267-277.
- Eskildsen, S., Hartmann, R., Kjeldgaard, N.O. and Justesen, J. 2002. Gene structure of the murine 2'- 5'- oligoadenylate synthetase family. *Cell. Mol. Life Sci.* 59: 1212-1222.
- Kakuta, S., Shibata, S. and Iwakura, Y. 2002. Genomic structure of the mouse 2'- 5'- oligoadenylate synthetase gene family. *J. Interferon Cytokine Res.* 22: 981-993.
- Eskildsen, S., Justesen, J., Schierup, M.H. and Hartmann, R. 2003. Characterization of the 2'- 5'- oligoadenylate synthetase ubiquitin-like family. *Nucleic Acids Res.* 31: 3166-3173.
- Andersen, J.B., Strandbygaard, D.J., Hartmann, R. and Justesen, J. 2004. Interaction between the 2'- 5'- oligoadenylate synthetase-like protein p59 OASL and the transcriptional repressor methyl CpG-binding protein 1. *Eur. J. Biochem.* 271: 628-636.

CHROMOSOMAL LOCATION

Genetic locus: OAS3 (human) mapping to 12q24.2; Oas3 (mouse) mapping to 5 F.

SOURCE

OAS3 (M-14) is an affinity purified goat polyclonal antibody raised against a peptide mapping at the C-terminus of OAS3 of mouse origin.

RESEARCH USE

For research use only, not for use in diagnostic procedures.

PRODUCT

Each vial contains 200 μ g IgG in 1.0 ml of PBS with < 0.1% sodium azide and 0.1% gelatin. Also available as TransCruz reagent for Gel Supershift and ChIP applications, sc-49869 X, 200 μ g/0.1 ml.

Blocking peptide available for competition studies, sc-49869 P, (100 μ g peptide in 0.5 ml PBS containing < 0.1% sodium azide and 0.2% BSA).

APPLICATIONS

OAS3 (M-14) is recommended for detection of OAS3 of mouse and rat origin by Western Blotting (starting dilution 1:200, dilution range 1:100-1:1000), immunofluorescence (starting dilution 1:50, dilution range 1:50-1:500) and solid phase ELISA (starting dilution 1:30, dilution range 1:30-1:3000).

Suitable for use as control antibody for OAS3 siRNA (m): sc-61246, OAS3 shRNA Plasmid (m): sc-61246-SH and OAS3 shRNA (m) Lentiviral Particles: sc-61246-V.

OAS3 (M-14) X TransCruz antibody is recommended for Gel Supershift and ChIP applications.

Molecular Weight of OAS3: 100 kDa.

RECOMMENDED SECONDARY REAGENTS

To ensure optimal results, the following support (secondary) reagents are recommended: 1) Western Blotting: use donkey anti-goat IgG-HRP: sc-2020 (dilution range: 1:2000-1:100,000) or Cruz Marker™ compatible donkey anti-goat IgG-HRP: sc-2033 (dilution range: 1:2000-1:5000), Cruz Marker™ Molecular Weight Standards: sc-2035, TBS Blotto A Blocking Reagent: sc-2333 and Western Blotting Luminol Reagent: sc-2048. 2) Immunofluorescence: use donkey anti-goat IgG-FITC: sc-2024 (dilution range: 1:100-1:400) or donkey anti-goat IgG-TR: sc-2783 (dilution range: 1:100-1:400) with UltraCruz™ Mounting Medium: sc-24941.

STORAGE

Store at 4° C, ****DO NOT FREEZE****. Stable for one year from the date of shipment. Non-hazardous. No MSDS required.

PROTOCOLS

See our web site at www.scbt.com or our catalog for detailed protocols and support products.



Try **OAS3 (D-7): sc-398225**, our highly recommended monoclonal alternative to OAS3 (M-14).