

HSP 40 (G-12): sc-49988

BACKGROUND

Heat shock protein 40 (HSP 40) family proteins bind to HSP 70 through their J-domain and regulate the function of HSP 70 by stimulating HSP 70 ATPase activity. HSP 40, also known as DnaJ, functions together with DnaK (HSP 70) and GrpE as a molecular chaperone, involving them in assembly and disassembly of protein complexes, protein folding, renaturation of denatured proteins, prevention of protein aggregation and protein export. HSP 40 stimulates the association between HSC 70 and HIP and translocates rapidly from the cytoplasm to the nuclei, and especially to the nucleoli, upon heat shock. There are five known HSP 40 family proteins.

REFERENCES

- Ohtsuka, K., Utsumi, K.R., Kaneda, T. and Hattori, H. 1993. Effect of ATP on the release of HSP 70 and HSP 40 from the nucleus in heat-shocked HeLa cells. *Exp. Cell Res.* 209: 357-366.
- Yamane, M., Hattori, H., Sugito, K., Hayash, Y., Tohnai, I., Ueda, M., Nishizawa, K. and Ohtsuka, K. 1995. Cotranslocation and co-localization of HSP 40 (DnaJ) with HSP 70 (DnaK) in mammalian cells. *Cell Struct. Funct.* 20: 157-166.
- Kaneko, R., Hattori, H., Hayashi, Y., Tohnai, I., Ueda, M. and Ohtsuka, K. 1995. Heat shock protein 40, a novel predictor of thermotolerance in murine cells. *Radiat. Res.* 142: 91-97.
- Hoe, K.L., 1998. Isolation of a new member of DnaJ-like heat shock protein 40 (HSP 40) from human liver. *Biochim. Biophys. Acta* 1383: 4-8.
- Michels, A.A., Kanon, B., Bensaude, O. and Kampinga, H.H. 1999. Heat shock protein (HSP) 40 mutants inhibit HSP 70 in mammalian cells. *J. Biol. Chem.* 274: 36757-36763.
- Suzuki, T., Usuda, N., Murata, S., Nakazawa, A., Ohtsuka, K. and Takagi, H. 1999. Presence of molecular chaperones, heat shock cognate (HSC) 70 and heat shock proteins (HSP) 40, in the postsynaptic structures of rat brain. *Brain Res.* 16: 816: 99-110.
- Ohtsuka, K. 2000. Mammalian HSP 40/DnaJ homologs: cloning of novel cDNAs and a proposal for their classification and nomenclature. *Cell Stress Chaperones* 5: 98-112.

CHROMOSOMAL LOCATION

Genetic locus: DNAJB1 (human) mapping to 19p13.12; Dnajb1 (mouse) mapping to 8 C2.

SOURCE

HSP 40 (G-12) is an affinity purified goat polyclonal antibody raised against a peptide mapping at the N-terminus of HSP 40 protein 1 of human origin.

PRODUCT

Each vial contains 200 µg IgG in 1.0 ml of PBS with < 0.1% sodium azide and 0.1% gelatin.

Blocking peptide available for competition studies, sc-49988 P, (100 µg peptide in 0.5 ml PBS containing < 0.1% sodium azide and 0.2% BSA).

APPLICATIONS

HSP 40 (G-12) is recommended for detection of HSP 40 protein 1 of mouse, rat and human origin by Western Blotting (starting dilution 1:200, dilution range 1:100-1:1000), immunofluorescence (starting dilution 1:50, dilution range 1:50-1:500) and solid phase ELISA (starting dilution 1:30, dilution range 1:30-1:3000).

HSP 40 (G-12) is also recommended for detection of HSP 40 protein 1 in additional species, including canine, bovine and porcine.

Suitable for use as control antibody for HSP 40 siRNA (h): sc-35599, HSP 40 siRNA (m): sc-40656, HSP 40 shRNA Plasmid (h): sc-35599-SH, HSP 40 shRNA Plasmid (m): sc-40656-SH, HSP 40 shRNA (h) Lentiviral Particles: sc-35599-V and HSP 40 shRNA (m) Lentiviral Particles: sc-40656-V.

Molecular Weight of HSP 40: 40 kDa.

Positive Controls: HeLa whole cell lysate: sc-2200, HeLa + heat shock cell lysate: sc-2272 or Hep G2 cell lysate: sc-2227.

RECOMMENDED SECONDARY REAGENTS

To ensure optimal results, the following support (secondary) reagents are recommended: 1) Western Blotting: use donkey anti-goat IgG-HRP: sc-2020 (dilution range: 1:2000-1:100,000) or Cruz Marker™ compatible donkey anti-goat IgG-HRP: sc-2033 (dilution range: 1:2000-1:5000), Cruz Marker™ Molecular Weight Standards: sc-2035, TBS Blotto A Blocking Reagent: sc-2333 and Western Blotting Luminol Reagent: sc-2048. 2) Immunofluorescence: use donkey anti-goat IgG-FITC: sc-2024 (dilution range: 1:100-1:400) or donkey anti-goat IgG-TR: sc-2783 (dilution range: 1:100-1:400) with UltraCruz™ Mounting Medium: sc-24941.

STORAGE

Store at 4° C, **DO NOT FREEZE**. Stable for one year from the date of shipment. Non-hazardous. No MSDS required.

RESEARCH USE

For research use only, not for use in diagnostic procedures.

PROTOCOLS

See our web site at www.scbt.com or our catalog for detailed protocols and support products.



Try **HSP 40 (B-3): sc-398766** or **HSP 40 (5): sc-135943**, our highly recommended monoclonal alternatives to HSP 40 (G-12).