

MYBPC3 (C-17): sc-50114

BACKGROUND

MYBPC3 encodes the cardiac isoform of the thick-filament myosin-binding protein C. It is found in the crossbridge-bearing zone (C region) of A bands in vertebrate striated muscle. Regulatory phosphorylation of MYBPC3 by cAMP-dependent protein kinase (PKA) upon adrenergic stimulation may be linked to modulation of cardiac contraction. MYBPC3 binds F-Actin, MHC and native thin filaments, and modifies the activity of Actin-activated myosin ATPase. Mutations in the MYBPC3 gene lead mainly to truncation of the protein, which results in one cause of familial hypertrophic cardiomyopathy type 4 (CMH4), a heart disorder characterized by ventricular hypertrophy, which often involves the interventricular septum and is usually asymmetric. The MYBPC3 gene maps to chromosome 11p11.2.

CHROMOSOMAL LOCATION

Genetic locus: MYBPC3 (human) mapping to 11p11.2; Mybpc3 (mouse) mapping to 2 E1.

SOURCE

MYBPC3 (C-17) is an affinity purified goat polyclonal antibody raised against a peptide mapping near the C-terminus of MYBPC3 of human origin.

PRODUCT

Each vial contains 200 µg IgG in 1.0 ml of PBS with < 0.1% sodium azide and 0.1% gelatin.

Blocking peptide available for competition studies, sc-50114 P, (100 µg peptide in 0.5 ml PBS containing < 0.1% sodium azide and 0.2% BSA).

STORAGE

Store at 4° C, ****DO NOT FREEZE****. Stable for one year from the date of shipment. Non-hazardous. No MSDS required.

APPLICATIONS

MYBPC3 (C-17) is recommended for detection of MYBPC3 of mouse, rat and human origin by Western Blotting (starting dilution 1:200, dilution range 1:100-1:1000), immunofluorescence (starting dilution 1:50, dilution range 1:50-1:500), immunohistochemistry (including paraffin-embedded sections) (starting dilution 1:50, dilution range 1:50-1:500) and solid phase ELISA (starting dilution 1:30, dilution range 1:30-1:3000).

MYBPC3 (C-17) is also recommended for detection of MYBPC3 in additional species, including canine, bovine and porcine.

Suitable for use as control antibody for MYBPC3 siRNA (h): sc-61111, MYBPC3 siRNA (m): sc-61112, MYBPC3 shRNA Plasmid (h): sc-61111-SH, MYBPC3 shRNA Plasmid (m): sc-61112-SH, MYBPC3 shRNA (h) Lentiviral Particles: sc-61111-V and MYBPC3 shRNA (m) Lentiviral Particles: sc-61112-V.


Molecular Weight of MYBPC3: 144 kDa.

Positive Controls: mouse heart extract: sc-2254 or rat heart extract: sc-2393.

RECOMMENDED SECONDARY REAGENTS

To ensure optimal results, the following support (secondary) reagents are recommended: 1) Western Blotting: use donkey anti-goat IgG-HRP: sc-2020 (dilution range: 1:2000-1:100,000) or Cruz Marker™ compatible donkey anti-goat IgG-HRP: sc-2033 (dilution range: 1:2000-1:5000), Cruz Marker™ Molecular Weight Standards: sc-2035, TBS Blotto A Blocking Reagent: sc-2333 and Western Blotting Luminol Reagent: sc-2048. 2) Immunofluorescence: use donkey anti-goat IgG-FITC: sc-2024 (dilution range: 1:100-1:400) or donkey anti-goat IgG-TR: sc-2783 (dilution range: 1:100-1:400) with UltraCruz™ Mounting Medium: sc-24941. 3) Immunohistochemistry: use ImmunoCruz™: sc-2053 or ABC: sc-2023 goat IgG Staining Systems.

DATA



MYBPC3 (C-17): sc-50114. Immunoperoxidase staining of formalin fixed, paraffin-embedded human heart muscle tissue showing cytoplasmic staining of myocytes.

RESEARCH USE

For research use only, not for use in diagnostic procedures.

PROTOCOLS

See our web site at www.scbt.com or our catalog for detailed protocols and support products.



Try **MYBPC3 (G-7): sc-137237** or **MYBPC3 (G-1): sc-137182**, our highly recommended monoclonal alternatives to MYBPC3 (C-17).