

neuroigin X/Y (D-20): sc-50135

BACKGROUND

Neuroigins are a family of plasma membrane proteins that possess an N-terminal hydrophobic sequence, a large esterase homology domain, a single transmembrane region and a short cytoplasmic domain. Neuroigins are expressed in excitatory, synaptic clefts and play a role in the formation and remodeling of CNS synapses by binding to β -neurexins, a family of neuronal cell surface proteins. Neuroigins also bind to PSD-95, which may recruit ion channels and neurotransmitter receptors to the synapse. A single-copy X-degenerate gene within the male-specific region of chromosome Y (MSY), which is not involved in X-Y crossover events, encodes neuroigin X/Y (NLGN4Y). Neuroigin X/Y and its X-linked homolog, NLGN4, contain all amino acids essential for neuroigin function, including the cysteines, the transmembrane domain and the PDZ-binding domain. Expression of NLGN4 and neuroigin X/Y is observed in all male brain regions examined, while female brain tissue only demonstrates NLGN4 expression.

REFERENCES

1. Nagase, T., Ishikawa, K., Suyama, M., Kikuno, R., Hirose, M., Miyajima, N., Tanaka, A., Kotani, H., Nomura, N. and Ohara, O. 1999. Prediction of the coding sequences of unidentified human genes. XIII. The complete sequences of 100 new cDNA clones from brain which code for large proteins *in vitro*. DNA Res. 6: 63-70.
2. Jamain, S., Quach, H., Betancur, C., Rastam, M., Colineaux, C., Gillberg, I.C., Soderstrom, H., Giros, B., Leboyer, M., Gillberg, C., Bourgeron, T. and Paris Autism Research International Sibpair Study. 2003. Mutations of the X-linked genes encoding neuroigins NLGN3 and NLGN4 are associated with autism. Nat. Genet. 34: 27-29.
3. Skaletsky, H., Kuroda-Kawaguchi, T., Minx, P.J., Cordum, H.S., Hillier, L., Brown, L.G., Repping, S., Pyntikova, T., Ali, J., Bieri, T., Chinwalla, A., Delehaunty, A., Delehaunty, K., Du, H., Fewell, G., Fulton, L., Fulton, R., et al. 2003. The male-specific region of the human Y chromosome is a mosaic of discrete sequence classes. Nature 423: 825-837.
4. Laumonnier, F., Bonnet-Brihault, F., Gomot, M., Blanc, R., David, A., Moizard, M.P., Raynaud, M., Ronce, N., Lemonnier, E., Calvas, P., Laudier, B., Chelly, J., Fryns, J.P., Ropers, H.H., Hamel, B.C. andres, C., Barthelemy, C., Moraine, C. and Briault, S. 2004. X-linked mental retardation and autism are associated with a mutation in the NLGN4 gene, a member of the neuroigin family. Am. J. Hum. Genet. 74: 552-557.
5. Vincent, J.B., Kolozsvari, D., Roberts, W.S., Bolton, P.F., Gurling, H.M. and Scherer, S.W. 2004. Mutation screening of X-chromosomal neuroigin genes: no mutations in 196 autism probands. Am. J. Med. Genet. B Neuropsychiatr. Genet. 129: 82-84.
6. Zhang, Z. and Henzel, W.J. 2004. Signal peptide prediction based on analysis of sites. Protein Sci. 13: 2819-2824.
7. Ylisaukko-oja, T., Rehnström, K., Auranen, M., Vanhala, R., Alen, R., Kempas, E., Ellonen, P., Turunen, J.A., Makkonen, I., Riikonen, R., Nieminen-von Wendt, T., von Wendt, L., Peltonen, L. and Järvelä, I. 2005. Analysis of four neuroigin genes as candidates for autism. Eur. J. Hum. Genet. 13: 1285-1292.

CHROMOSOMAL LOCATION

Genetic locus: NLGN4X (human) mapping to Xp22.32, NLGN4Y (human) mapping to Yq11.221.

SOURCE

neuroigin X/Y (D-20) is an affinity purified goat polyclonal antibody raised against a peptide mapping within an extracellular domain of neuroigin X/Y of human origin.

PRODUCT

Each vial contains 200 μ g IgG in 1.0 ml of PBS with < 0.1% sodium azide and 0.1% gelatin.

Blocking peptide available for competition studies, sc-50135 P, (100 μ g peptide in 0.5 ml PBS containing < 0.1% sodium azide and 0.2% BSA).

APPLICATIONS

neuroigin X/Y (D-20) is recommended for detection of neuroigin X/Y of human origin by Western Blotting (starting dilution 1:200, dilution range 1:100-1:1000), immunofluorescence (starting dilution 1:50, dilution range 1:50-1:500) and solid phase ELISA (starting dilution 1:30, dilution range 1:30-1:3000).

neuroigin X/Y (D-20) is also recommended for detection of neuroigin X/Y in additional species, including equine, canine, bovine and porcine.

Suitable for use as control antibody for neuroigin X/Y siRNA (h): sc-61186, neuroigin X/Y shRNA Plasmid (h): sc-61186-SH and neuroigin X/Y shRNA (h) Lentiviral Particles: sc-61186-V.

Molecular Weight of neuroigin X/Y: 92 kDa.

RECOMMENDED SECONDARY REAGENTS

To ensure optimal results, the following support (secondary) reagents are recommended: 1) Western Blotting: use donkey anti-goat IgG-HRP: sc-2020 (dilution range: 1:2000-1:100,000) or Cruz Marker™ compatible donkey anti-goat IgG-HRP: sc-2033 (dilution range: 1:2000-1:5000), Cruz Marker™ Molecular Weight Standards: sc-2035, TBS Blotto A Blocking Reagent: sc-2333 and Western Blotting Luminol Reagent: sc-2048. 2) Immunofluorescence: use donkey anti-goat IgG-FITC: sc-2024 (dilution range: 1:100-1:400) or donkey anti-goat IgG-TR: sc-2783 (dilution range: 1:100-1:400) with UltraCruz™ Mounting Medium: sc-24941.

STORAGE

Store at 4° C, **DO NOT FREEZE**. Stable for one year from the date of shipment. Non-hazardous. No MSDS required.

RESEARCH USE

For research use only, not for use in diagnostic procedures.

PROTOCOLS

See our web site at www.scbt.com or our catalog for detailed protocols and support products.