

XK (W-13): sc-50201



The Power to Question

BACKGROUND

Kell and XK are two covalently linked plasma membrane proteins that constitute the Kell blood group system, a group of antigens on the surface of red blood cells that are important determinants of blood type and targets for autoimmune or alloimmune diseases. XK is a 444 amino acid protein that spans the membrane 10 times and carries the ubiquitous antigen, Kx, which determines blood type. XK also plays a role in the sodium-dependent membrane transport of oligopeptides and neutral amino acids. XK is expressed at high levels in brain, heart, skeletal muscle and pancreas. Defects in the XK gene cause McLeod syndrome (MLS), an X-linked multisystem disorder characterized by abnormalities in neuromuscular and hematopoietic system such as acanthocytic red blood cells and late-onset forms of muscular dystrophy with nerve abnormalities.

REFERENCES

1. Lee, S., Russo, D. and Redman, C. 2000. Functional and structural aspects of the Kell blood group system. *Transfus. Med. Rev.* 14: 93-103.
2. Lee, S., Russo, D. and Redman, C.M. 2000. The Kell blood group system: Kell and XK membrane proteins. *Semin. Hematol.* 37: 113-121.
3. Online Mendelian Inheritance in Man, OMIM™. 2002. Johns Hopkins University, Baltimore, MD. MIM Number: 314850. World Wide Web URL: <http://www.ncbi.nlm.nih.gov/omim/>
4. Singleton, B.K., Green, C.A., Renaud, S., Fuhr, P., Poole, J. and Daniels, G.L. 2003. McLeod syndrome resulting from a novel XK mutation. *Br. J. Haematol.* 122: 682-685.
5. Danek, A., Jung, H.H., Melone, M.A., Rampoldi, L., Broccoli, V. and Walker, R.H. 2005. Neuroacanthocytosis: new developments in a neglected group of dementing disorders. *J. Neurol. Sci.* 229-230: 171-186.
6. Feder, M. and Bujnicki, J.M. 2005. Identification of a new family of putative PD-(D/E)XK nucleases with unusual phylogenomic distribution and a new type of the active site. *BMC Genomics* 6: 21.
7. Pu, J.J., Redman, C.M., Visser, J.W. and Lee, S. 2005. Onset of expression of the components of the Kell blood group complex. *Transfusion* 45: 969-974.
8. Starling, A., Schlesinger, D., Kok, F., Passos-Bueno, M.R., Vainzof, M. and Zatz, M. 2005. A family with McLeod syndrome and calpainopathy with clinically overlapping diseases. *Neurology* 65: 1832-1833.
9. Zeman, A., Daniels, G., Tilley, L., Dunn, M., Toplis, L., Bullock, T., Poole, J. and Blackwood, D. 2005. McLeod syndrome: life-long neuropsychiatric disorder due to a novel mutation of the XK gene. *Psychiatr. Genet.* 15: 291-293.

CHROMOSOMAL LOCATION

Genetic locus: XK (human) mapping to Xp21.1; Xk (mouse) mapping to X A1.1.

SOURCE

XK (W-13) is an affinity purified goat polyclonal antibody raised against a peptide mapping within an internal region of XK of human origin.

PRODUCT

Each vial contains 200 µg IgG in 1.0 ml of PBS with < 0.1% sodium azide and 0.1% gelatin.

Blocking peptide available for competition studies, sc-50201 P, (100 µg peptide in 0.5 ml PBS containing < 0.1% sodium azide and 0.2% BSA).

APPLICATIONS

XK (W-13) is recommended for detection of XK of mouse, rat and human origin by Western Blotting (starting dilution 1:200, dilution range 1:100-1:1000), immunofluorescence (starting dilution 1:50, dilution range 1:50-1:500) and solid phase ELISA (starting dilution 1:30, dilution range 1:30-1:3000).

XK (W-13) is also recommended for detection of XK in additional species, including equine, canine, bovine and porcine.

Suitable for use as control antibody for XK siRNA (h): sc-61807, XK siRNA (m): sc-61808, XK shRNA Plasmid (h): sc-61807-SH, XK shRNA Plasmid (m): sc-61808-SH, XK shRNA (h) Lentiviral Particles: sc-61807-V and XK shRNA (m) Lentiviral Particles: sc-61808-V.

Molecular Weight of XK: 40 kDa.

RECOMMENDED SECONDARY REAGENTS

To ensure optimal results, the following support (secondary) reagents are recommended: 1) Western Blotting: use donkey anti-goat IgG-HRP: sc-2020 (dilution range: 1:2000-1:100,000) or Cruz Marker™ compatible donkey anti-goat IgG-HRP: sc-2033 (dilution range: 1:2000-1:5000), Cruz Marker™ Molecular Weight Standards: sc-2035, TBS Blotto A Blocking Reagent: sc-2333 and Western Blotting Luminol Reagent: sc-2048. 2) Immunofluorescence: use donkey anti-goat IgG-FITC: sc-2024 (dilution range: 1:100-1:400) or donkey anti-goat IgG-TR: sc-2783 (dilution range: 1:100-1:400) with UltraCruz™ Mounting Medium: sc-24941.

STORAGE

Store at 4° C, **DO NOT FREEZE**. Stable for one year from the date of shipment. Non-hazardous. No MSDS required.

RESEARCH USE

For research use only, not for use in diagnostic procedures.

PROTOCOLS

See our web site at www.scbt.com or our catalog for detailed protocols and support products.