

CKIP-1 (S-20): sc-50227

BACKGROUND

Casein kinase II-interacting protein 1 (CKIP-1), also designated Pleckstrin homology domain containing family O member 1 (PLEKH01), is a 409 amino acid protein with an N-terminal Pleckstrin homology domain and a putative C-terminal Jun leucine zipper interactive domain. CKIP-1 is expressed at the highest levels in skeletal muscle and heart, intermediately in placenta, lung and brain and at the weakest levels in pancreas, liver and kidney. CKIP-1 localizes to the plasma membrane of transfected COS-7 cells and also to the plasma membrane and the nucleus in human osteosarcoma cells. It interacts with the N-terminus of CSNK2A1 and with full length CSNK2A1, but not with CSNK2A2 or CSNK2B.

REFERENCES

1. Bosc, D.G., Graham, K.C., Saulnier, R.B., Zhang, C., Prober, D., Gietz, R.D. and Litchfield, D.W. 2000. Identification and characterization of CKIP-1, a novel Pleckstrin homology domain-containing protein that interacts with protein kinase CK2. *J. Biol. Chem.* 275: 14295-14306.
2. Olsten, M.E., Canton, D.A., Zhang, C., Walton, P.A. and Litchfield, D.W. 2004. The Pleckstrin homology domain of CK2 interacting protein-1 is required for interactions and recruitment of protein kinase CK2 to the plasma membrane. *J. Biol. Chem.* 279: 42114-42127.
3. Barrios-Rodiles, M., Brown, K.R., Ozdamar, B., Bose, R., Liu, Z., Donovan, R.S., Shinjo, F., Liu, Y., Dembowy, J., Taylor, I.W., Luga, V., Przulj, N., Robinson, M., Suzuki, H., Hayashizaki, Y., Jurisica, I. and Wrana, J.L. 2005. High-throughput mapping of a dynamic signaling network in mammalian cells. *Science* 307: 1621-1625.
4. Canton, D.A., Olsten, M.E., Kim, K., Doherty-Kirby, A., Lajoie, G., Cooper, J.A. and Litchfield, D.W. 2005. The Pleckstrin homology domain-containing protein CKIP-1 is involved in regulation of cell morphology and the Actin cytoskeleton and interaction with Actin capping protein. *Mol. Cell. Biol.* 25: 3519-3534.

CHROMOSOMAL LOCATION

Genetic locus: PLEKH01 (human) mapping to 1q21.2; Plekho1 (mouse) mapping to 3 F2.

SOURCE

CKIP-1 (S-20) is an affinity purified goat polyclonal antibody raised against a peptide mapping within an internal region of CKIP-1 of human origin.

PRODUCT

Each vial contains 200 µg IgG in 1.0 ml of PBS with < 0.1% sodium azide and 0.1% gelatin.

Blocking peptide available for competition studies, sc-50227 P, (100 µg peptide in 0.5 ml PBS containing < 0.1% sodium azide and 0.2% BSA).

RESEARCH USE

For research use only, not for use in diagnostic procedures.

APPLICATIONS

CKIP-1 (S-20) is recommended for detection of CKIP-1 of mouse, rat and human origin by Western Blotting (starting dilution 1:200, dilution range 1:100-1:1000), immunoprecipitation [1-2 µg per 100-500 µg of total protein (1 ml of cell lysate)], immunofluorescence (starting dilution 1:50, dilution range 1:50-1:500) and solid phase ELISA (starting dilution 1:30, dilution range 1:30-1:3000).

Suitable for use as control antibody for CKIP-1 siRNA (h): sc-60389, CKIP-1 siRNA (m): sc-60390, CKIP-1 shRNA Plasmid (h): sc-60389-SH, CKIP-1 shRNA Plasmid (m): sc-60390-SH, CKIP-1 shRNA (h) Lentiviral Particles: sc-60389-V and CKIP-1 shRNA (m) Lentiviral Particles: sc-60390-V.


Molecular Weight of CKIP-1: 50 kDa.

Positive Controls: CKIP-1 (h4): 293T Lysate: sc-176668.

RECOMMENDED SECONDARY REAGENTS

To ensure optimal results, the following support (secondary) reagents are recommended: 1) Western Blotting: use donkey anti-goat IgG-HRP: sc-2020 (dilution range: 1:2000-1:100,000) or Cruz Marker™ compatible donkey anti-goat IgG-HRP: sc-2033 (dilution range: 1:2000-1:5000), Cruz Marker™ Molecular Weight Standards: sc-2035, TBS Blotto A Blocking Reagent: sc-2333 and Western Blotting Luminol Reagent: sc-2048. 2) Immunoprecipitation: use Protein A/G PLUS-Agarose: sc-2003 (0.5 ml agarose/2.0 ml). 3) Immunofluorescence: use donkey anti-goat IgG-FITC: sc-2024 (dilution range: 1:100-1:400) or donkey anti-goat IgG-TR: sc-2783 (dilution range: 1:100-1:400) with UltraCruz™ Mounting Medium: sc-24941.

DATA



CKIP-1 (S-20): sc-50227. Western blot analysis of CKIP-1 expression in non-transfected: sc-117752 (A) and human CKIP-1 transfected: sc-176668 (B) 293T whole cell lysates.

STORAGE

Store at 4° C, **DO NOT FREEZE**. Stable for one year from the date of shipment. Non-hazardous. No MSDS required.

MONOS
Satisfaction
Guaranteed

Try **CKIP-1 (A-12): sc-376355** or **CKIP-1 (A-3): sc-376060**, our highly recommended monoclonal alternatives to CKIP-1 (S-20).