

c-Fgr (M-60): sc-50338

BACKGROUND

Src is the human homolog of the v-Src gene of the rous sarcoma virus, also designated avian sarcoma virus or ASV. Src was the first proto-oncogenic non-receptor tyrosine kinase characterized in human. The Src family, which has common structural motifs, is composed of nine members in vertebrates, including Src, Yes, Fgr, Frk, Fyn, Lyn, Hck, Lck and Blk. Src-family kinases transduce signals that are involved in the control of a variety of cellular processes, including proliferation, differentiation, motility and adhesion. Src-family kinases contain an amino-terminal cell membrane anchor followed by an SH3 domain and an SH2 domain, which are involved in modular association and activation, respectively. Src-family kinases, which are normally maintained in an inactive state and can be activated transiently during cellular events such as mitosis. Different subcellular localizations of Src-family kinases may be important for the regulation of specific cellular processes such as mitogenesis, cytoskeletal organization and membrane trafficking. c-Fgr is a human non-receptor tyrosine kinase family member that was discovered by using a probe toward the v-Fgr portion of the cell-derived domain of Gardner-Rasheed feline sarcoma virus. The human c-Fgr gene encodes a 529 amino acid protein.

REFERENCES

1. Sakaguchi, A.Y., et al. 1982. Organization of human proto-oncogenes. *Am. J. Hum. Gen.* 34: 175.
2. Tronick, S.R., et al. 1985. Isolation and chromosomal localization of the human Fgr protooncogene, a distinct member of the tyrosine kinase gene family. *Proc. Natl. Acad. Sci. USA* 82: 6595-6599.
3. Williams, J.C., et al. 1998. Insights into Src kinase functions: structural comparisons. *Trends Biochem. Sci.* 23: 179-184.
4. Tatosyan, A.G., et al. 2000. Kinases of the Src-family: structure and functions. *Biochemistry* 65: 49-58.
5. Bjorge, J.D., et al. 2000. Selected glimpses into the activation and function of Src kinase. *Oncogene* 19: 5620-5635.
6. Online Mendelian Inheritance in Man, OMIM[™]. 2000. Johns Hopkins University, Baltimore, MD. MIM Number: 190090. World Wide Web URL: <http://www.ncbi.nlm.nih.gov/omim/>

CHROMOSOMAL LOCATION

Genetic locus: FGR (human) mapping to 1p36.11; Fgr (mouse) mapping to 4 D2.3.

SOURCE

c-Fgr (M-60) is a rabbit polyclonal antibody raised against amino acids 4-63 mapping at the N-terminus of c-Fgr of mouse origin.

PRODUCT

Each vial contains 200 µg IgG in 1.0 ml of PBS with < 0.1% sodium azide and 0.1% gelatin.

RESEARCH USE

For research use only, not for use in diagnostic procedures.

APPLICATIONS

c-Fgr (M-60) is recommended for detection of c-Fgr p55 of mouse, rat and human origin by Western Blotting (starting dilution 1:200, dilution range 1:100-1:1000), immunoprecipitation [1-2 µg per 100-500 µg of total protein (1 ml of cell lysate)], immunofluorescence (starting dilution 1:50, dilution range 1:50-1:500) and solid phase ELISA (starting dilution 1:30, dilution range 1:30-1:3000).

Suitable for use as control antibody for c-Fgr siRNA (h): sc-39229, c-Fgr siRNA (m): sc-39230, c-Fgr shRNA Plasmid (h): sc-39229-SH, c-Fgr shRNA Plasmid (m): sc-39230-SH, c-Fgr shRNA (h) Lentiviral Particles: sc-39229-V and c-Fgr shRNA (m) Lentiviral Particles: sc-39230-V.


Molecular Weight of c-Fgr: 55 kDa.

Positive Controls: RAW 264.7 whole cell lysate: sc-2211 or HeLa whole cell lysate: sc-2200.


RECOMMENDED SECONDARY REAGENTS

To ensure optimal results, the following support (secondary) reagents are recommended: 1) Western Blotting: use goat anti-rabbit IgG-HRP: sc-2004 (dilution range: 1:2000-1:100,000) or Cruz Marker[™] compatible goat anti-rabbit IgG-HRP: sc-2030 (dilution range: 1:2000-1:5000), Cruz Marker[™] Molecular Weight Standards: sc-2035, TBS Blotto A Blocking Reagent: sc-2333 and Western Blotting Luminol Reagent: sc-2048. 2) Immunoprecipitation: use Protein A/G PLUS-Agarose: sc-2003 (0.5 ml agarose/2.0 ml). 3) Immunofluorescence: use goat anti-rabbit IgG-FITC: sc-2012 (dilution range: 1:100-1:400) or goat anti-rabbit IgG-TR: sc-2780 (dilution range: 1:100-1:400) with UltraCruz[™] Mounting Medium: sc-24941.

DATA



c-Fgr (M-60): sc-50338. Western blot analysis of c-Fgr expression in RAW 264.7 whole cell lysate.



c-Fgr (M-60): sc-50338. Western blot analysis of human recombinant c-Fgr fusion protein.

STORAGE

Store at 4° C, ****DO NOT FREEZE****. Stable for one year from the date of shipment. Non-hazardous. No MSDS required.



Try **c-Fgr (B-8): sc-166079** or **c-Fgr (D-6): sc-74542**, our highly recommended monoclonal alternatives to c-Fgr (M-60).