

## zero (H-60): sc-50406

### BACKGROUND

Zero, also known as myelin protein zero (MPZ) is a Type 1 integral membrane glycoprotein that mediates adhesion of spiraling wraps of the myelin sheath in order to ensure stable synaptic transmission. Zero protein encompasses approximately 50% of total protein in the sheath scaffolding in contribution to structural integrity of peripheral myelin. Zero guides the compact myelin wrapping process through glycine zipper packing interface-dependent dimer and tetramer formation. Mutations (e.g. G134R) can abrogate multimer formation, cause demyelinating neuropathies, and are known to contribute to conditions that include Charcot-Marie-Tooth disease. Zero cytoplasmic domain undergoes serine and tyrosine phosphorylation, which appears to be prevalent during peak nerve myelination. Zero transcript is moderate in brain, abundant in thymus and most abundant in white matter of the CNS.

### REFERENCES

1. Shy, M.E., Jáni, A., Krajewski, K., Grandis, M., Lewis, R.A., Li, J., Shy, R.R., Balsamo, J., Lilien, J., Garbern, J.Y. and Kamholz, J. 2004. Phenotypic clustering in MPZ mutations. *Brain* 127: 371-384.
2. Plotkowski, M.L., Kim, S., Phillips, M.L., Partridge, A.W., Deber, C.M. and Bowie, J.U. 2007. Transmembrane domain of myelin protein zero can form dimers: possible implications for myelin construction. *Biochemistry* 46: 12164-12173.
3. Gaboreanu, A.M., Hrstka, R., Xu, W., Shy, M., Kamholz, J., Lilien, J. and Balsamo, J. 2007. Myelin protein zero/P0 phosphorylation and function require an adaptor protein linking it to RACK1 and PKC  $\alpha$ . *J. Cell Biol.* 177: 707-716.
4. Taguchi, K., Kumanogoh, H., Nakamura, S., Miyata, S. and Maekawa, S. 2007. Myelin protein zero is one of the components of the detergent-resistant membrane microdomain fraction prepared from rat pituitary. *J. Mol. Histol.* 38: 79-85.
5. Kim, H.J., Jung, C.G., Jensen, M.A., Dukala, D. and Soliven, B. 2008. Targeting of myelin protein zero in a spontaneous autoimmune polyneuropathy. *J. Immunol.* 181: 8753-8760.
6. Lemke, G. 2008. Isolation and analysis of the gene encoding peripheral myelin protein zero. *Neuron* 60: 403.

### CHROMOSOMAL LOCATION

Genetic locus: MPZ (human) mapping to 1q23.3; Mpz (mouse) mapping to 1 H3.

### SOURCE

zero (H-60) is a rabbit polyclonal antibody raised against amino acids 11-70 mapping within an extracellular domain of zero of human origin.

### PRODUCT

Each vial contains 200  $\mu$ g IgG in 1.0 ml of PBS with < 0.1% sodium azide and 0.1% gelatin.

### APPLICATIONS

zero (H-60) is recommended for detection of zero of mouse, rat and human origin by Western Blotting (starting dilution 1:200, dilution range 1:100-1:1000), immunoprecipitation [1-2  $\mu$ g per 100-500  $\mu$ g of total protein (1 ml of cell lysate)], immunofluorescence (starting dilution 1:50, dilution range 1:50-1:500) and solid phase ELISA (starting dilution 1:30, dilution range 1:30-1:3000).

zero (H-60) is also recommended for detection of zero in additional species, including porcine.

Suitable for use as control antibody for zero siRNA (h): sc-44194, zero siRNA (m): sc-44497, zero shRNA Plasmid (h): sc-44194-SH, zero shRNA Plasmid (m): sc-44497-SH, zero shRNA (h) Lentiviral Particles: sc-44194-V and zero shRNA (m) Lentiviral Particles: sc-44497-V.


Molecular Weight of zero: 28 kDa.

Positive Controls: human spinal cord tissue extract.

### RECOMMENDED SECONDARY REAGENTS

To ensure optimal results, the following support (secondary) reagents are recommended: 1) Western Blotting: use goat anti-rabbit IgG-HRP: sc-2004 (dilution range: 1:2000-1:100,000) or Cruz Marker™ compatible goat anti-rabbit IgG-HRP: sc-2030 (dilution range: 1:2000-1:5000), Cruz Marker™ Molecular Weight Standards: sc-2035, TBS Blotto A Blocking Reagent: sc-2333 and Western Blotting Luminol Reagent: sc-2048. 2) Immunoprecipitation: use Protein A/G PLUS-Agarose: sc-2003 (0.5 ml agarose/2.0 ml). 3) Immunofluorescence: use goat anti-rabbit IgG-FITC: sc-2012 (dilution range: 1:100-1:400) or goat anti-rabbit IgG-TR: sc-2780 (dilution range: 1:100-1:400) with UltraCruz™ Mounting Medium: sc-24941.

### DATA



zero (H-60): sc-50406. Western blot analysis of zero expression in human spinal cord tissue extract.

### STORAGE

Store at 4° C, \*\*DO NOT FREEZE\*\*. Stable for one year from the date of shipment. Non-hazardous. No MSDS required.

### RESEARCH USE

For research use only, not for use in diagnostic procedures.

### PROTOCOLS

See our web site at [www.scbt.com](http://www.scbt.com) or our catalog for detailed protocols and support products.