

# Myo-inositol oxygenase (N-19): sc-50603

## BACKGROUND

Myo-inositol oxygenase (MIOX), also known as ALDRL6, is a renal-specific member of the Aldo-keto reductase family. It catalyzes the first committed step in the Myo-inositol metabolism pathway and is widely distributed in mammalian tissues. Human Myo-inositol oxygenase shares 91% and 96% sequence homology with mouse and pig Myo-inositol oxygenase homologs, respectively. Myo-inositol oxygenase is responsible for the oxidative cleavage of Myo-inositol (MI) and its epimer D-chiro inositol (DCI) to D-glucuronate. The dioxygen-dependent cleavage of the C1-C6 bond in Myo-inositol is accomplished through the utilization of the Fe(II)/Fe(III) binuclear iron center of MIOX. Myo-inositol oxygenase has also been implicated in complications of diabetes, including diabetic nephropathy.

## CHROMOSOMAL LOCATION

Genetic locus: MIOX (human) mapping to 22q13.33; Miox (mouse) mapping to 15 E3.

## SOURCE

Myo-inositol oxygenase (N-19) is an affinity purified goat polyclonal antibody raised against a peptide mapping at the N-terminus of Myo-inositol oxygenase of human origin.

## PRODUCT

Each vial contains 200 µg IgG in 1.0 ml of PBS with < 0.1% sodium azide and 0.1% gelatin.

Blocking peptide available for competition studies, sc-50603 P, (100 µg peptide in 0.5 ml PBS containing < 0.1% sodium azide and 0.2% BSA).

## APPLICATIONS

Myo-inositol oxygenase (N-19) is recommended for detection of Myo-inositol oxygenase of mouse, rat and human origin by Western Blotting (starting dilution 1:200, dilution range 1:100-1:1000), immunoprecipitation (1-2 µg per 100-500 µg of total protein (1 ml of cell lysate)), immunofluorescence (starting dilution 1:50, dilution range 1:50-1:500) and solid phase ELISA (starting dilution 1:30, dilution range 1:30-1:3000).

Myo-inositol oxygenase (N-19) is also recommended for detection of Myo-inositol oxygenase in additional species, including equine and canine.

Suitable for use as control antibody for Myo-inositol oxygenase siRNA (h): sc-61117, Myo-inositol oxygenase siRNA (m): sc-61118, Myo-inositol oxygenase shRNA Plasmid (h): sc-61117-SH, Myo-inositol oxygenase shRNA Plasmid (m): sc-61118-SH, Myo-inositol oxygenase shRNA (h) Lentiviral Particles: sc-61117-V and Myo-inositol oxygenase shRNA (m) Lentiviral Particles: sc-61118-V.

Molecular Weight of Myo-inositol oxygenase: 33 kDa.

Positive Controls: Myo-inositol oxygenase (h2): 293T Lysate: sc-117367, mouse kidney extract: sc-2255 or rat kidney extract: sc-2394.


## RESEARCH USE

For research use only, not for use in diagnostic procedures.


## STORAGE

Store at 4° C, **\*\*DO NOT FREEZE\*\***. Stable for one year from the date of shipment. Non-hazardous. No MSDS required.

## DATA



Myo-inositol oxygenase (N-19): sc-50603. Western blot analysis of Myo-inositol oxygenase expression in human kidney (A), mouse kidney (B) and rat kidney (C) tissue extracts.



Myo-inositol oxygenase (N-19): sc-50603. Western blot analysis of Myo-inositol oxygenase expression in non-transfected: sc-117752 (A) and human Myo-inositol oxygenase transfected: sc-117367 (B) 293T whole cell lysates and mouse kidney tissue extract (C).

## PROTOCOLS

See our web site at [www.scbt.com](http://www.scbt.com) or our catalog for detailed protocols and support products.



**MONOS**  
Satisfation  
Guaranteed

Try **Myo-inositol oxygenase (E-11): sc-376080** or **Myo-inositol oxygenase (E-9): sc-166913**, our highly recommended monoclonal alternatives to Myo-inositol oxygenase (N-19).