

ASCT2 (A-14): sc-50695

BACKGROUND

Neutral amino acid transporter proteins, also designated alanine/serine/cysteine/threonine transporters (ASCT), belong to the sodium:dicarboxylate (SDF) symporter family of proteins. The members of this family of proteins are multi-pass membrane-bound proteins that act as transporters for threonine, alanine, serine and cysteine. ASCT1 and ASCT2 have been shown to exhibit sodium dependence. ASCT1 is expressed in most tissues, but highest expression has been detected in muscle, brain and pancreas. The highest levels of ASCT2 expression are found in placenta, kidney, pancreas, muscle and intestine.

REFERENCES

1. Arriza, J.L., Kavanaugh, M.P., Fairman, W.A., Wu, Y.N., Murdoch, G.H., North, R.A. and Amara, S.G. 1993. Cloning and expression of a human neutral amino acid transporter with structural similarity to the glutamate transporter gene family. *J. Biol. Chem.* 268: 15329-15332.
2. Hofmann, K., Düker, M., Fink, T., Lichter, P. and Stoffel, W. 1995. Human neutral amino acid transporter ASCT1: structure of the gene (SLC1A4) and localization to chromosome 2p13-p15. *Genomics* 24: 20-26.
3. Kekuda, R., Prasad, P.D., Fei, Y.J., Torres-Zamorano, V., Sinha, S., Yang-Feng, T.L., Leibach, F.H. and Ganapathy, V. 1996. Cloning of the sodium-dependent, broad-scope, neutral amino acid transporter Bo from a human placental choriocarcinoma cell line. *J. Biol. Chem.* 271: 18657-18661.
4. Tailor, C.S., Nouri, A., Zhao, Y., Takeuchi, Y. and Kabat, D. 1999. A sodium-dependent neutral-amino-acid transporter mediates infections of feline and baboon endogenous retroviruses and simian type D retroviruses. *J. Virol.* 73: 4470-4474.
5. Rasko, J.E., Battini, J.L., Gottschalk, R.J., Mazo, I. and Miller, A.D. 1999. The RD114/simian type D retrovirus receptor is a neutral amino acid transporter. *Proc. Natl. Acad. Sci. USA* 96: 2129-2134.
6. Tailor, C.S., Marin, M., Nouri, A., Kavanaugh, M.P. and Kabat, D. 2001. Truncated forms of the dual function human ASCT2 neutral amino acid transporter/retroviral receptor are translationally initiated at multiple alternative CUG and GUG codons. *J. Biol. Chem.* 276: 27221-27230.
7. Kanai, Y. and Hediger, M.A. 2004. The glutamate/neutral amino acid transporter family SLC1: molecular, physiological and pharmacological aspects. *Pflügers Archiv.* 447: 469-79.
8. Yamamoto, T., Nishizaki, I., Nukada, T., Kamegaya, E., Furuya, S., Hirabayashi, Y., Ikeda, K., Hata, H., Kobayashi, H., Sora, I. and Yamamoto, H. 2004. Functional identification of ASCT1 neutral amino acid transporter as the predominant system for the uptake of L-serine in rat neurons in primary culture. *Neurosci. Res.* 49: 101-111.
9. Hashimoto, Y., Sadamoto, Y., Konno, A., Kon, Y. and Iwanaga, T. 2005. Distribution of neutral amino acid transporter ASCT 1 in the non-neuronal tissues of mice. *Jpn. J. Vet. Res.* 52: 113-124.

CHROMOSOMAL LOCATION

Genetic locus: Slc1a5 (mouse) mapping to 7 A2.

SOURCE

ASCT2 (A-14) is an affinity purified goat polyclonal antibody raised against a peptide mapping within an extracellular domain of ASCT2 of mouse origin.

PRODUCT

Each vial contains 200 µg IgG in 1.0 ml of PBS with < 0.1% sodium azide and 0.1% gelatin.

Blocking peptide available for competition studies, sc-50695 P, (100 µg peptide in 0.5 ml PBS containing < 0.1% sodium azide and 0.2% BSA).

APPLICATIONS

ASCT2 (A-14) is recommended for detection of ASCT2 of mouse origin by Western Blotting (starting dilution 1:200, dilution range 1:100-1:1000), immunofluorescence (starting dilution 1:50, dilution range 1:50-1:500) and solid phase ELISA (starting dilution 1:30, dilution range 1:30-1:3000).

Suitable for use as control antibody for ASCT2 siRNA (m): sc-60211, ASCT2 shRNA Plasmid (m): sc-60211-SH and ASCT2 shRNA (m) Lentiviral Particles: sc-60211-V.

Molecular Weight of endogenous ASCT2: 55 kDa.

Molecular Weight of glycosylated ASCT2: 70-80 kDa.

RECOMMENDED SECONDARY REAGENTS

To ensure optimal results, the following support (secondary) reagents are recommended: 1) Western Blotting: use donkey anti-goat IgG-HRP: sc-2020 (dilution range: 1:2000-1:100,000) or Cruz Marker™ compatible donkey anti-goat IgG-HRP: sc-2033 (dilution range: 1:2000-1:5000), Cruz Marker™ Molecular Weight Standards: sc-2035, TBS Blotto A Blocking Reagent: sc-2333 and Western Blotting Luminol Reagent: sc-2048. 2) Immunofluorescence: use donkey anti-goat IgG-FITC: sc-2024 (dilution range: 1:100-1:400) or donkey anti-goat IgG-TR: sc-2783 (dilution range: 1:100-1:400) with UltraCruz™ Mounting Medium: sc-24941.

STORAGE

Store at 4° C, **DO NOT FREEZE**. Stable for one year from the date of shipment. Non-hazardous. No MSDS required.

RESEARCH USE

For research use only, not for use in diagnostic procedures.

PROTOCOLS

See our web site at www.scbt.com or our catalog for detailed protocols and support products.