

# Glyoxalase I (T-16): sc-50732

## BACKGROUND

The glyoxal pathway plays a role in the detoxification of glucose degradation products (GDP). Glyoxalase I (GLO1), a member of the Glyoxalase family, is effective in eliminating GDP. Overexpression or silencing of Glyoxalase I in mouse brain suggests an association between Glyoxalase I and anxiety. Glyoxalase I has three isoforms generated from two alleles in the genome which forms two homodimers and one heterodimer, each subunit binding one zinc ion. Research demonstrates that GLO1 gene expression is induced in colon carcinoma. Both an Insulin response element (IRE) and a zinc metal response element (MRE) in the promoter region of the GLO1 gene have been identified.

## CHROMOSOMAL LOCATION

Genetic locus: GLO1 (human) mapping to 6p21.2; Glo1 (mouse) mapping to 17 A3.3.

## SOURCE

Glyoxalase I (T-16) is an affinity purified goat polyclonal antibody raised against a peptide mapping within an internal region of Glyoxalase I of human origin.

## PRODUCT

Each vial contains 200 µg IgG in 1.0 ml of PBS with < 0.1% sodium azide and 0.1% gelatin.

Blocking peptide available for competition studies, sc-50732 P, (100 µg peptide in 0.5 ml PBS containing < 0.1% sodium azide and 0.2% BSA).

## STORAGE

Store at 4° C, **\*\*DO NOT FREEZE\*\***. Stable for one year from the date of shipment. Non-hazardous. No MSDS required.

## APPLICATIONS

Glyoxalase I (T-16) is recommended for detection of Glyoxalase I of mouse, rat and human origin by Western Blotting (starting dilution 1:200, dilution range 1:100-1:1000), immunoprecipitation [1-2 µg per 100-500 µg of total protein (1 ml of cell lysate)], immunofluorescence (starting dilution 1:50, dilution range 1:50-1:500) and solid phase ELISA (starting dilution 1:30, dilution range 1:30-1:3000).

Glyoxalase I (T-16) is also recommended for detection of Glyoxalase I in additional species, including equine, canine, bovine and porcine.

Suitable for use as control antibody for Glyoxalase I siRNA (h): sc-60703, Glyoxalase I siRNA (m): sc-60704, Glyoxalase I shRNA Plasmid (h): sc-60703-SH, Glyoxalase I shRNA Plasmid (m): sc-60704-SH, Glyoxalase I shRNA (h) Lentiviral Particles: sc-60703-V and Glyoxalase I shRNA (m) Lentiviral Particles: sc-60704-V.


Molecular Weight of Glyoxalase I monomer: 24 kDa.

Positive Controls: HEL 92.1.7 cell lysate: sc-2270, TF-1 cell lysate: sc-2412 or HeLa whole cell lysate: sc-2200.


## RECOMMENDED SECONDARY REAGENTS

To ensure optimal results, the following support (secondary) reagents are recommended: 1) Western Blotting: use donkey anti-goat IgG-HRP: sc-2020 (dilution range: 1:2000-1:100,000) or Cruz Marker™ compatible donkey anti-goat IgG-HRP: sc-2033 (dilution range: 1:2000-1:5000), Cruz Marker™ Molecular Weight Standards: sc-2035, TBS Blotto A Blocking Reagent: sc-2333 and Western Blotting Luminol Reagent: sc-2048. 2) Immunoprecipitation: use Protein A/G PLUS-Agarose: sc-2003 (0.5 ml agarose/2.0 ml). 3) Immunofluorescence: use donkey anti-goat IgG-FITC: sc-2024 (dilution range: 1:100-1:400) or donkey anti-goat IgG-TR: sc-2783 (dilution range: 1:100-1:400) with UltraCruz™ Mounting Medium: sc-24941.

## DATA



Glyoxalase I (T-16): sc-50732. Western blot analysis of Glyoxalase I expression in HEL 92.1.7 (A) and TF-1 (B) whole cell lysates.



Glyoxalase I (T-16): sc-50732. Immunofluorescence staining of methanol-fixed HeLa cells showing cytoplasmic localization.

## RESEARCH USE

For research use only, not for use in diagnostic procedures.

## PROTOCOLS

See our web site at [www.scbt.com](http://www.scbt.com) or our catalog for detailed protocols and support products.


 MONOS  
Satisfaction  
Guaranteed

Try **Glyoxalase I (D-5): sc-133214** or **Glyoxalase I (D-6): sc-133144**, our highly recommended monoclonal alternatives to Glyoxalase I (T-16).