AHSP (W-20): sc-50795



The Power to Question

BACKGROUND

 α -hemoglobin stabilizing protein (AHSP), also designated erythroid associated factor (ERAF), is an erythroid-specific protein that acts as a chaperone to prevent the aggregation of α -hemoglobin during normal erythroid cell development. It specifically protects free α -hemoglobin from precipitation in live cells and in solution. It forms a heterodimer with free α -hemoglobin, but not with β -hemoglobin or hemoglobin A (α 2- β -2). AHSP localizes to the cytoplasm and is expressed in the blood and bone marrow. The AHSP protein is down-regulated in transmissible spongiform encephalopathies (TSEs). AHSP may regulate pathological states of α -hemoglobin excess such as β -thalassemia, a group of hereditary disorders involving the decreased production of normal adult hemoglobin (HbA) that are characterized by a deficiency in the synthesis of β -globin chains.

REFERENCES

- Zhang, Q.H., Ye, M., Wu, X.Y., Ren, S.X., Zhao, M., Zhao, C.J., Fu, G., Shen, Y., Fan, H.Y., Lu, G., Zhong, M., Xu, X.R., Han, Z.G., Zhang, J.W., Tao, J., Huang, Q.H., Zhou, J., Hu, G.X., Gu, J., Chen, S.J. and Chen, Z. 2000. Cloning and functional analysis of cDNAs with open reading frames for 300 previously undefined genes expressed in CD34+ hematopoietic stem/ progenitor cells. Genome Res. 10: 1546-1560.
- Miele, G., Manson, J. and Clinton, M. 2001. A novel erythroid-specific marker of transmissible spongiform encephalopathies. Nat. Med. 7: 361-364.
- Kihm, A.J., Kong, Y., Hong, W., Russell, J.E., Rouda, S., Adachi, K., Simon, M.C., Blobel, G.A. and Weiss, M.J. 2002. An abundant erythroid protein that stabilizes free α-haemoglobin. Nature 417: 758-763.
- 4. Gell, D., Kong, Y., Eaton, S.A., Weiss, M.J. and Mackay, J.P. 2002. Biophysical characterization of the α -globin binding protein α -hemoglobin stabilizing protein. J. Biol. Chem. 277: 40602-40609.
- 5. Online Mendelian Inheritance in Man, OMIM™. 2002. Johns Hopkins University, Baltimore, MD. MIM Number: 605821. World Wide Web URL: http://www.ncbi.nlm.nih.gov/omim/
- 6. dos Santos, C.O. and Costa, F.F. 2005. AHSP and β-thalassemia: a possible genetic modifier. Hematology 10: 157-161.

SOURCE

AHSP (W-20) is an affinity purified goat polyclonal antibody raised against a peptide mapping near the N-terminus of AHSP of mouse origin.

PRODUCT

Each vial contains 200 μg lgG in 1.0 ml of PBS with < 0.1% sodium azide and 0.1% gelatin.

Blocking peptide available for competition studies, sc-50795 P, (100 μg peptide in 0.5 ml PBS containing < 0.1% sodium azide and 0.2% BSA).

STORAGE

Store at 4° C, **DO NOT FREEZE**. Stable for one year from the date of shipment. Non-hazardous. No MSDS required.

APPLICATIONS

AHSP (W-20) is recommended for detection of AHSP of mouse and rat origin by Western Blotting (starting dilution 1:200, dilution range 1:100-1:1000), immunoprecipitation [1-2 μ g per 100-500 μ g of total protein (1 ml of cell lysate)], immunofluorescence (starting dilution 1:50, dilution range 1:50-1:500) and solid phase ELISA (starting dilution 1:30, dilution range 1:30-1:3000).

Suitable for use as control antibody for AHSP siRNA (m): sc-60138, AHSP shRNA Plasmid (m): sc-60138-SH and AHSP shRNA (m) Lentiviral Particles: sc-60138-V.

Molecular Weight of AHSP: 12 kDa.

RECOMMENDED SECONDARY REAGENTS

To ensure optimal results, the following support (secondary) reagents are recommended: 1) Western Blotting: use donkey anti-goat IgG-HRP: sc-2020 (dilution range: 1:2000-1:100,000) or Cruz Marker™ compatible donkey anti-goat IgG-HRP: sc-2033 (dilution range: 1:2000-1:5000), Cruz Marker™ Molecular Weight Standards: sc-2035, TBS Blotto A Blocking Reagent: sc-2333 and Western Blotting Luminol Reagent: sc-2048. 2) Immunoprecipitation: use Protein A/G PLUS-Agarose: sc-2003 (0.5 ml agarose/2.0 ml). 3) Immunofluorescence: use donkey anti-goat IgG-FITC: sc-2024 (dilution range: 1:100-1:400) or donkey anti-goat IgG-TR: sc-2783 (dilution range: 1:100-1:400) with UltraCruz™ Mounting Medium: sc-24941.

RESEARCH USE

For research use only, not for use in diagnostic procedures.

PROTOCOLS

See our web site at www.scbt.com or our catalog for detailed protocols and support products.



Try **AHSP (G-5): sc-515189**, our highly recommended monoclonal alternative to AHSP (W-20).

Santa Cruz Biotechnology, Inc. 1.800.457.3801 831.457.3801 Fax 831.457.3801 Europe +00800 4573 8000 49 6221 4503 0 www.scbt.com