# MASP-1 (L-17): sc-50840



The Power to Question

## **BACKGROUND**

Mannose (or mannan)-binding lectin (MBL), also known as serum mannosebinding protein (MBP), initiates the lectin branch of the innate immune response by binding to the surface of potentially pathogenic microorganisms and initiating complement fixation through an N-terminal collagen-like domain. MBL is a key component in immune response in that it can directly trigger neutralization of invading microorganisms by an Ab-independent mechanism. Mutations of human MBL are associated with immunodeficiency resulting from a reduction in the ability of the mutant MBL to initiate complement fixation. In human, three types of MBL-associated serine proteases, MASP-1, MASP-2 and MASP-3, and a truncated form of MASP-2 (small MBL-associated protein; sMAP or MAp19) complex with MBL to activate the lectin pathway of the complement system. MASP-3 is an alternatively spliced product from the MASP-1 gene. The heavy/A chains are identical between MASP-1 and MASP-3 but the light/B chains are entirely different. Activated MASPs subsequently cleave and activate downstream components of the complement pathway.

## **REFERENCES**

- Heise, C., et al. 2000. Impaired secretion of rat mannose-binding protein resulting from mutations in the collagen-like domain. J. Immunol. 165: 1403-1409.
- Matsushita, M., et al. 2000. Proteolytic activities of two types of mannose-binding lectin-associated serine protease. J. Immunol. 165: 2637-2642.
- Chen, C.B., et al. 2001. Stoichiometry of complexes between mannosebinding protein and its associated serine proteases: Defining functional units for complement activation. J. Biol. Chem. 276: 25894-25902.
- 4. Endo, M., et al. 2001. Regulation of *in situ* complement activation via the lectin pathway in patients with IgA nephropathy. Clin. Nephrol. 55: 185-191.
- 5. Thielens, N.M., et al. 2001. Interaction properties of human mannan-binding lectin (MBL)-associated serine proteases-1 and -2, MBL-associated protein 19, and MBL. J. Immunol. 166: 5068-5077.

#### CHROMOSOMAL LOCATION

Genetic locus: Masp1 (mouse) mapping to 16 B1.

#### **SOURCE**

MASP-1 (L-17) is an affinity purified goat polyclonal antibody raised against a peptide mapping within an internal region of the processed light chain region of MASP-1 of mouse origin.

#### **PRODUCT**

Each vial contains 200  $\mu g$  lgG in 1.0 ml of PBS with < 0.1% sodium azide and 0.1% gelatin.

Blocking peptide available for competition studies, sc-50840 P, (100  $\mu$ g peptide in 0.5 ml PBS containing < 0.1% sodium azide and 0.2% BSA).

#### **APPLICATIONS**

MASP-1 (L-17) is recommended for detection of MASP-1 light chain region of mouse and rat origin by Western Blotting (starting dilution 1:200, dilution range 1:100-1:1000), immunofluorescence (starting dilution 1:50, dilution range 1:50-1:500) and solid phase ELISA (starting dilution 1:30, dilution range 1:30-1:3000).

MASP-1 (L-17) is also recommended for detection of MASP-1 light chain region in additional species, including equine and canine.

Suitable for use as control antibody for MASP-1 siRNA (m): sc-45350, MASP-1 shRNA Plasmid (m): sc-45350-SH and MASP-1 shRNA (m) Lentiviral Particles: sc-45350-V.

Molecular Weight of MASP-1: 35 kDa.

## **RECOMMENDED SECONDARY REAGENTS**

To ensure optimal results, the following support (secondary) reagents are recommended: 1) Western Blotting: use donkey anti-goat IgG-HRP: sc-2020 (dilution range: 1:2000-1:100,000) or Cruz Marker™ compatible donkey anti-goat IgG-HRP: sc-2033 (dilution range: 1:2000-1:5000), Cruz Marker™ Molecular Weight Standards: sc-2035, TBS Blotto A Blocking Reagent: sc-2333 and Western Blotting Luminol Reagent: sc-2048. 2) Immunofluorescence: use donkey anti-goat IgG-FITC: sc-2024 (dilution range: 1:100-1:400) or donkey anti-goat IgG-TR: sc-2783 (dilution range: 1:100-1:400) with UltraCruz™ Mounting Medium: sc-24941.

## **STORAGE**

Store at 4° C, \*\*DO NOT FREEZE\*\*. Stable for one year from the date of shipment. Non-hazardous. No MSDS required.

## **RESEARCH USE**

For research use only, not for use in diagnostic procedures.

### **PROTOCOLS**

See our web site at www.scbt.com or our catalog for detailed protocols and support products.

Santa Cruz Biotechnology, Inc. 1.800.457.3801 831.457.3801 Fax 831.457.3801 Europe +00800 4573 8000 49 6221 4503 0 www.scbt.com