SANTA CRUZ BIOTECHNOLOGY, INC.

CA I (C-20): sc-50895



BACKGROUND

Carbonic anhydrases (CAs), also designated carbonate dehydratases or carbonate hydrolyases, form a large family of genes that encode zinc metalloenzymes of great physiologic importance. As catalysts of the reversible hydration of carbon dioxide, these enzymes participate in a variety of biologic processes, including respiration, acid-base balance, bone resorption and calcification, as well as the formation of aqueous humor, cerebrospinal fluid, saliva and gastric acid. Genes in the α -carbonic anhydrase family encode either active carbonic anhydrase isozymes or "acatalytic" (devoid of CO₂ hydration activity) carbonic anhydrase-related proteins. Human CA I (CA1) is encoded by the CA1 gene, which has been assigned to chromosome 8 and harbors a cluster of CA genes. CA I localizes to the cytoplasm and research indicates that a severe deficiency of CA I does not result in any obvious hematological or renal consequences.

REFERENCES

- Hopkinson, D.A., et al. 1975. The detection and differentiation of the products of the human carbonic anhydrase loci, CA I and CA II using fluorogenic substrates. Ann. Hum. Genet. 38: 155-162.
- Davis, M.B., et al. 1987. Regional localization of carbonic anhydrase genes CA1 and CA3 on human chromosome 8. Somat. Cell Mol. Genet. 13: 173-178.
- Edwards, Y.H., et al. 1988. Assignment of the gene determining human carbonic anhydrase, CA1, to chromosome 8. Ann. Hum. Genet. 50 (Pt 2): 123-129.
- 4. Hewett-Emmett, D. and Tashian, R.E. 1996. Functional diversity, conservation, and convergence in the evolution of the α -, β -, and γ -carbonic anhydrase gene families. Mol. Phylogenet. Evol. 5: 50-77.
- Ichihara, N., et al. 1998. Immunohistolocalization of carbonic anhydrase isozymes (CA I, CA II and CA III) in bovine male reproductive tracts. Okajimas Folia Anat. Jpn. 74: 193-198.

CHROMOSOMAL LOCATION

Genetic locus: CA1 (human) mapping to 8q21.2; Car1 (mouse) mapping to 3 A1.

SOURCE

CA I (C-20) is an affinity purified goat polyclonal antibody raised against a peptide mapping at the C-terminus of CA I of human origin.

PRODUCT

Each vial contains 200 μg IgG in 1.0 ml of PBS with < 0.1% sodium azide and 0.1% gelatin.

Blocking peptide available for competition studies, sc-50895 P, (100 μ g peptide in 0.5 ml PBS containing < 0.1% sodium azide and 0.2% BSA).

STORAGE

Store at 4° C, **DO NOT FREEZE**. Stable for one year from the date of shipment. Non-hazardous. No MSDS required.

APPLICATIONS

CA I (C-20) is recommended for detection of CA I of human, and, to a lesser extent, mouse origin by Western Blotting (starting dilution 1:200, dilution range 1:100-1:1000), immunoprecipitation [1-2 µg per 100-500 µg of total protein (1 ml of cell lysate)], immunofluorescence (starting dilution 1:50, dilution range 1:50-1:500) and solid phase ELISA (starting dilution 1:30, dilution range 1:30-1:3000).

CA I (C-20) is also recommended for detection of CA I (carbonic anhydrase) in additional species, including canine.

Molecular Weight of CA I: 29 kDa.

Positive Controls: K-562 whole cell lysate: sc-2203, HEL 92.1.7 cell lysate: sc-2270 or CA I (h): 293T Lysate: sc-114146.

RECOMMENDED SECONDARY REAGENTS

To ensure optimal results, the following support (secondary) reagents are recommended: 1) Western Blotting: use donkey anti-goat IgG-HRP: sc-2020 (dilution range: 1:2000-1:100,000) or Cruz Marker™ compatible donkey anti-goat IgG-HRP: sc-2033 (dilution range: 1:2000-1:5000), Cruz Marker™ Molecular Weight Standards: sc-2035, TBS Blotto A Blocking Reagent: sc-2333 and Western Blotting Luminol Reagent: sc-2048. 2) Immunoprecipitation: use Protein A/G PLUS-Agarose: sc-2003 (0.5 ml agarose/2.0 ml). 3) Immunofluorescence: use donkey anti-goat IgG-FITC: sc-2024 (dilution range: 1:100-1:400) or donkey anti-goat IgG-TR: sc-2783 (dilution range: 1:100-1:400) with UltraCruz™ Mounting Medium: sc-24941.

DATA





CA I (C-20): sc-50895. Western blot analysis of CA I expression in non-transfected: sc-117752 (A) and human CA I transfected: sc-114146 (B) 293T whole cell lysates.

CA I (C-20): sc-50895. Western blot analysis of CA I expression in mouse spleen tissue extract.

RESEARCH USE

For research use only, not for use in diagnostic procedures.

MONOS Satisfation Guaranteed Try CA I (F-5): sc-393490 or CA I (F-11): sc-393497, our highly recommended monoclonal alternatives to CA I (C-20).