

## PFK-2 car (N-20): sc-50953

### BACKGROUND

Phosphofructokinases (PFK) are regulatory glycolytic enzymes that convert fructose 6-phosphate and ATP into fructose 1,6-bisphosphate (through PFK-1), fructose 2,6-bisphosphate (through PFK-2) and ADP. Human PFK-1 is tetrameric; isoenzymes include PFK-1 muscle (PFKM, PFK-A), PFK-1 liver (PFKL, PFK-B) and PFK-1 platelet (PFKP, PFK-C, PFKF). PFK-1 is inhibited by ATP and citrate (from the tricarboxylic acid cycle). PFK-1 undergoes activation in the presence of elevated AMP. The most potent activator is fructose-2,6-bisphosphate, which is produced by PFK-2 from the same substrate, fructose 6-phosphate. PFK-2 is bifunctional and a key regulator for PFK-1. PFK-2 catalyzes the synthesis of fructose-2,6-bisphosphate, and contains fructose-2,6-bisphosphatase activity that catalyzes the degradation of fructose-2,6-bisphosphate. PFK-2 is dimeric; isoenzymes include PFK-2 liver (PFKFB1, PFRX), PFK-2 cardiac (PFKFB2), PFK-2 placental (PFKFB3, inducible PFK-2) and PFK-2 testis (PFKFB4).

### REFERENCES

1. Tsuura, Y., et al. 1998. Endogenous nitric oxide inhibits glucose-induced Insulin secretion by suppression of phosphofructokinase activity in pancreatic islets. *Biochem. Biophys. Res. Commun.* 252: 34-38.
2. Chang, S.H., et al. 2002. Role of Ser 530, Arg 292, and His 662 in the allosteric behavior of rabbit muscle phosphofructokinase. *Biochem. Biophys. Res. Commun.* 290: 670-675.
3. Zeitschel, U., et al. 2002. Changes in activity and expression of phosphofructokinase in different rat brain regions after basal forebrain cholinergic lesion. *J. Neurochem.* 83: 371-380.
4. Su, Y., et al. 2003. The  $\alpha$  subunit of the V-type H<sup>+</sup>-ATPase interacts with phosphofructokinase-1 in humans. *J. Biol. Chem.* 278: 20013-20018.
5. Sotgia, F., et al. 2003. Phosphofructokinase muscle-specific isoform requires caveolin-3 expression for plasma membrane recruitment and caveolar targeting: implications for the pathogenesis of caveolin-related muscle diseases. *Am. J. Pathol.* 163: 2619-2634.
6. Haller, R.G., et al. 2004. No spontaneous second wind in muscle phosphofructokinase deficiency. *Neurology* 62: 82-86.
7. Martin, S.R., et al. 2004. Interaction of calmodulin with the phosphofructokinase target sequence. *FEBS Lett.* 577: 284-288.
8. Yamada, S., et al. 2004. Novel testis- and embryo-specific isoforms of the phosphofructokinase-1 muscle type gene. *Biochem. Biophys. Res. Commun.* 316: 580-587.
9. Vallejo, J., et al. 2005. Expression of caveolin-1 in lymphocytes induces caveolae formation and recruitment of phosphofructokinase to the plasma membrane. *FASEB J.* 19: 586-587.

### CHROMOSOMAL LOCATION

Genetic locus: PFKFB2 (human) mapping to 1q32.2.

### RESEARCH USE

For research use only, not for use in diagnostic procedures.

### SOURCE

PFK-2 car (N-20) is an affinity purified goat polyclonal antibody raised against a peptide mapping at the N-terminus of PFK-2 car of human origin.

### PRODUCT

Each vial contains 200  $\mu$ g IgG in 1.0 ml of PBS with < 0.1% sodium azide and 0.1% gelatin.

Blocking peptide available for competition studies, sc-50953 P, (100  $\mu$ g peptide in 0.5 ml PBS containing < 0.1% sodium azide and 0.2% BSA).

### APPLICATIONS

PFK-2 car (N-20) is recommended for detection of PFK-2 car of human origin by Western Blotting (starting dilution 1:200, dilution range 1:100-1:1000), immunofluorescence (starting dilution 1:50, dilution range 1:50-1:500) and solid phase ELISA (starting dilution 1:30, dilution range 1:30-1:3000).

PFK-2 car (N-20) is also recommended for detection of PFK-2 car in additional species, including canine and porcine.

Suitable for use as control antibody for PFK-2 car siRNA (h): sc-44675, PFK-2 car shRNA Plasmid (h): sc-44675-SH and PFK-2 car shRNA (h) Lentiviral Particles: sc-44675-V.

Molecular Weight of PFK-2 car: 58 kDa.

### RECOMMENDED SECONDARY REAGENTS

To ensure optimal results, the following support (secondary) reagents are recommended: 1) Western Blotting: use donkey anti-goat IgG-HRP: sc-2020 (dilution range: 1:2000-1:100,000) or Cruz Marker™ compatible donkey anti-goat IgG-HRP: sc-2033 (dilution range: 1:2000-1:5000), Cruz Marker™ Molecular Weight Standards: sc-2035, TBS Blotto A Blocking Reagent: sc-2333 and Western Blotting Luminol Reagent: sc-2048. 2) Immunofluorescence: use donkey anti-goat IgG-FITC: sc-2024 (dilution range: 1:100-1:400) or donkey anti-goat IgG-TR: sc-2783 (dilution range: 1:100-1:400) with UltraCruz™ Mounting Medium: sc-24941.

### STORAGE

Store at 4° C, \*\*DO NOT FREEZE\*\*. Stable for one year from the date of shipment. Non-hazardous. No MSDS required.

### PROTOCOLS

See our web site at [www.scbt.com](http://www.scbt.com) or our catalog for detailed protocols and support products.



Try **PFK-2 car (D-1): sc-377416** or **PFK-2 car (G-7): sc-515528**, our highly recommended monoclonal alternatives to PFK-2 car (N-20).